



Synonym

TNFRSF25,DR3,APO3,DDR3,TNFRSF12,WSL,WSL1

Source

Biotinylated Human DR3 Protein, Fc,Avitag(DR3-H82F3) is expressed from human 293 cells (HEK293). It contains AA Gln 25 - Gln 199 (Accession # [Q93038-1](#)).

Molecular Characterization

DR3(Gln 25 - Gln 199)	Fc(Pro 100 - Lys 330)	Avi
Q93038-1	P01857	

This protein carries a human IgG1 Fc tag at the C-terminus, followed by an Avi tag (Avitag™).

The protein has a calculated MW of 47.2 kDa. The protein migrates as 55-65 kDa when calibrated against [Star Ribbon Pre-stained Protein Marker](#) under reducing (R) condition (SDS-PAGE) due to glycosylation.

Labeling

Biotinylation of this product is performed using Avitag™ technology. Briefly, the single lysine residue in the Avitag is enzymatically labeled with biotin.

Protein Ratio

Passed as determined by the HABA assay / binding ELISA.

Purity

>90% as determined by SDS-PAGE.

>95% as determined by SEC-MALS.

Formulation

Lyophilized from 0.22 µm filtered solution in PBS, pH7.4 with trehalose as protectant.

Contact us for customized product form or formulation.

Reconstitution

Please see Certificate of Analysis for specific instructions.

For best performance, we strongly recommend you to follow the reconstitution protocol provided in the CoA.

Storage


For long term storage, the product should be stored at lyophilized state at -20°C or lower.

Please avoid repeated freeze-thaw cycles.

This product is stable after storage at:


- -20°C to -70°C for 12 months in lyophilized state;
- -70°C for 3 months under sterile conditions after reconstitution.

SDS-PAGE



Biotinylated Human DR3 Protein, Fc,Avitag on SDS-PAGE under reducing (R) condition. The gel was stained with Coomassie Blue. The purity of the protein is greater than 90% (With [Star Ribbon Pre-stained Protein Marker](#)).

SEC-MALS



The purity of Biotinylated Human DR3 Protein, Fc,Avitag (Cat. No. DR3-H82F3) is more than 95% and the molecular weight of this protein is around 95-120 kDa verified by SEC-MALS.


[Report](#)

Bioactivity-ELISA


Discounts, Gifts,
and more!




» www.acrobiosystems.com




Immobilized Human TL1A, His Tag (Cat. No. TLA-H5243) at 5 µg/mL (100 µL/well) can bind Biotinylated Human DR3 Protein, Fc,Avitag (Cat. No. DR3-H82F3) with a linear range of 0.005-0.625 µg/mL (QC tested).



Immobilized Human TL1A Protein, His Tag (Cat. No. TLA-H5244) at 5 µg/mL (100 µL/well) can bind Biotinylated Human DR3 Protein, Fc,Avitag (Cat. No. DR3-H82F3) with a linear range of 0.02-0.625 µg/mL (Routinely tested).




Immobilized Human TL1A, His Tag (Cat. No. TLA-H5243) at 2 µg/mL (100 µL/well) can bind pre-mixed increasing concentrations of Monoclonal Anti-TL1A Antibody, Human IgG1 and 4 µg/mL (50 µL/well) Biotinylated Human DR3 Protein, Fc,Avitag (Cat. No. DR3-H82F3) with a half maximal inhibitory concentration (IC50) of 1.976 µg/mL (Routinely tested).



Immobilized Human TL1A Protein, His Tag (Cat. No. TLA-H5244) at 2 µg/mL (100 µL/well) can bind pre-mixed increasing concentrations of Monoclonal Anti-TL1A Antibody, Human IgG1 and 4 µg/mL (50 µL/well) Biotinylated Human DR3 Protein, Fc,Avitag (Cat. No. DR3-H82F3) with a half maximal inhibitory concentration (IC50) of 3.06 µg/mL (Routinely tested).

Bioactivity-SPR



Biotinylated Human DR3 Protein, Fc,Avitag (Cat. No. DR3-H82F3) captured on Biotin CAP-Series S Sensor Chip can bind Human TL1A, His Tag (Cat. No.

Discounts, Gifts,
and more!





TLA-H5243) with an affinity constant of 1.02 μ M as determined in a SPR assay (Biacore 8K) (Routinely tested).

Background

Tumor necrosis factor receptor superfamily member 25 (TNFRSF25) is also known as Apo-3, Death receptor 3 (DDR3 or DR3), Apoptosis-inducing receptor AIR, Apoptosis-mediating receptor TRAMP, Lymphocyte-associated receptor of death, Apo-3, which is a member of the TNF-receptor superfamily. TNFRSF25 is a homodimer protein, which can interact strongly via the death domains with TNFRSF1 and TRADD to activate at least two distinct signaling cascades, apoptosis and NF- κ B signaling. TNFRSF25 is receptor for TNFSF12 / APO3L / TWEAK.

Discounts, Gifts,
and more!



» www.acrobiosystems.com