



Synonym

EGFR,ERBB,ERBB1,HER1,PIG61,mENA

Source

APC-Labeled Human EGF R Protein, His Tag (EGR-HA2H8) is produced via conjugation of APC to Human EGF R Protein, His Tag with a new generation site-specific technology under Star Staining labeling platform. Human EGF R Protein, His Tag is expressed from human 293 cells (HEK293). It contains AA Leu 25 - Ser 645 (Accession # [P00533-1](#)).

Predicted N-terminus: Leu 25

Molecular Characterization

This protein carries a polyhistidine tag at the C-terminus.

Conjugate

APC

Excitation Wavelength: 640 nm

Emission Wavelength: 661 nm

Purity

>90% as determined by SDS-PAGE.

Formulation

Lyophilized from 0.22 µm filtered solution in PBS, 0.2% BSA, pH7.4 with trehalose as protectant.

Contact us for customized product form or formulation.

Reconstitution

Please see Certificate of Analysis for specific instructions.

For best performance, we strongly recommend you to follow the reconstitution protocol provided in the CoA.

Storage

For long term storage, the product should be stored at lyophilized state at -20°C or lower.

Please protect from light and avoid repeated freeze-thaw cycles.

This product is stable after storage at:


- -20°C to -70°C for 12 months in lyophilized state;
- -70°C for 3 months under sterile conditions after reconstitution.

Star Staining fluorescent-labeled products are developed by a new-generation site-specific labeling technology with Star Standard quality at ACROBiosystems

- ★ Using new-generation site-specific labeling technology to maintain natural bioactivity.
- ★ No non-specific binding to non-transduced PBMCs.
- ★ High specificity and sensitivity verified by flow cytometry.
- ★ High homogeneity and high batch-to-batch consistency.

Evaluation of CAR expression

FACS Analysis of Anti-EGF R CAR Expression



Discounts, Gifts,
and more!




» www.acrobiosystems.com



5e5 of anti-EGFR CAR-293 cells were stained with 100 μ L of 1:50 dilution (2 μ L stock solution in 100 μ L FACS buffer) of APC-Labeled Human EGF R Protein, His Tag (Cat. No. EGR-HA2H8) and negative control protein respectively (Fig. C and B), and non-transfected 293 cells were used as a control (Fig. A). APC signal was used to evaluate the binding activity (QC tested).

FACS Analysis of Non-specific binding to PBMCs



5e5 of PBMCs were stained with APC-Labeled Human EGF R Protein, His Tag (Cat. No. EGR-HA2H8) and anti-CD3 antibody, washed and then analyzed with FACS. FITC signal was used to evaluate the expression of CD3+ T cells in PBMCs, and APC signal was used to evaluate the non-specific binding activity to PBMCs (QC tested).

Background

The epidermal growth factor receptor (EGFR; ErbB-1; HER1 in humans) is the cell-surface receptor for members of the epidermal growth factor family (EGF-family) of extracellular protein ligands. The epidermal growth factor receptor is a member of the ErbB family of receptors, a subfamily of four closely related receptor tyrosine kinases: EGFR (ErbB-1), HER2/c-neu (ErbB-2), Her 3 (ErbB-3) and Her 4 (ErbB-4). Mutations affecting EGFR expression or activity could result in cancer.

Discounts, Gifts,
and more!



» www.acrobiosystems.com

7/31/2025