

**Synonym**

EphA2

Source

Rat EphA2 Protein, His Tag(EP2-R52H5) is expressed from human 293 cells (HEK293). It contains AA Gln 24 - Met 536 (Accession # [D3ZBN3](#)).

Predicted N-terminus: Gln 24

Molecular Characterization

EphA2(Gln 24 - Met 536)	Poly-his
D3ZBN3	

This protein carries a polyhistidine tag at the C-terminus.

The protein has a calculated MW of 58.6 kDa. The protein migrates as 60-67 kDa when calibrated against [Star Ribbon Pre-stained Protein Marker](#) under reducing (R) condition (SDS-PAGE) due to glycosylation.

Endotoxin

Less than 1.0 EU per μ g by the LAL method / rFC method.

Purity

>95% as determined by SDS-PAGE.

>90% as determined by SEC-MALS.

Formulation

Lyophilized from 0.22 μ m filtered solution in PBS, pH7.4 with trehalose as protectant.

Contact us for customized product form or formulation.

Reconstitution

Please see Certificate of Analysis for specific instructions.

For best performance, we strongly recommend you to follow the reconstitution protocol provided in the CoA.


Storage

For long term storage, the product should be stored at lyophilized state at -20°C or lower.


Please avoid repeated freeze-thaw cycles.

This product is stable after storage at:

- -20°C to -70°C for 12 months in lyophilized state;
- -70°C for 3 months under sterile conditions after reconstitution.

SDS-PAGE

Rat EphA2 Protein, His Tag on SDS-PAGE under reducing (R) condition. The gel was stained with Coomassie Blue. The purity of the protein is greater than 95% (With [Star Ribbon Pre-stained Protein Marker](#)).

SEC-MALS

The purity of Rat EphA2 Protein, His Tag (Cat. No. EP2-R52H5) is more than 90% and the molecular weight of this protein is around 50-65 kDa verified by SEC-MALS.



[Report](#)

Bioactivity-ELISA

Discounts, Gifts,
and more!



» www.acrobiosystems.com



Immobilized Human Ephrin-A1, Fc Tag (Cat. No. EF1-H5251) at 5 µg/mL (100 µL/well) can bind Rat EphA2 Protein, His Tag (Cat. No. EP2-R52H5) with a linear range of 0.01-0.156 (QC tested).

Immobilized Rat EphA2 Protein, His Tag (Cat. No. EP2-R52H5) at 5 µg/mL (100 µL/well) can bind Human Ephrin-A1, Fc Tag (Cat. No. EF1-H5251) with a linear range of 0.04-1 ng/mL (Routinely tested).

Background

Receptor tyrosine kinase which binds promiscuously membrane-bound ephrin-A family ligands residing on adjacent cells, leading to contact-dependent bidirectional signaling into neighboring cells. The signaling pathway downstream of the receptor is referred to as forward signaling while the signaling pathway downstream of the ephrin ligand is referred to as reverse signaling. Activated by the ligand ephrin-A1/EF1 regulates migration, integrin-mediated adhesion, proliferation and differentiation of cells. Regulates cell adhesion and differentiation through DSG1/desmoglein-1 and inhibition of the ERK1/ERK2 (MAPK3/MAPK1, respectively) signaling pathway. Engaged by the ligand ephrin-A5/EF5 may regulate lens fiber cells shape and interactions and be important for lens transparency development and maintenance. With ephrin-A2/EF2 may play a role in bone remodeling through regulation of osteoclastogenesis and osteoblastogenesis.

Discounts, Gifts,
and more!

