

Synonym

FCGR1A,FCG1,FCGR1,IGFR1,CD64,CD64A,FCRI

Source

Canine CD64, His Tag(FCA-C52H8) is expressed from human 293 cells (HEK293). It contains AA Gln 16 - Pro 288 (Accession # [XP_025308990.1](#)).

Predicted N-terminus: Gln 16

Molecular Characterization

CD64(Gln 16 - Pro 288)
XP_025308990.1 Poly-his

This protein carries a polyhistidine tag at the C-terminus

The protein has a calculated MW of 32.4 kDa. The protein migrates as 36-46 kDa under reducing (R) condition (SDS-PAGE) due to glycosylation.

Endotoxin

Less than 1.0 EU per µg by the LAL method.

Purity

>90% as determined by SDS-PAGE.

>90% as determined by SEC-MALS.

Formulation

Lyophilized from 0.22 µm filtered solution in PBS, pH7.4 with Trehalose as protectant.

Contact us for customized product form or formulation.

Reconstitution

Please see Certificate of Analysis for specific instructions.

For best performance, we strongly recommend you to follow the reconstitution protocol provided in the CoA.


Storage

For long term storage, the product should be stored at lyophilized state at -20°C or lower.


Please avoid repeated freeze-thaw cycles.

This product is stable after storage at:

- -20°C to -70°C for 12 months in lyophilized state;
- -70°C for 3 months under sterile conditions after reconstitution.


SDS-PAGE

Canine CD64, His Tag on SDS-PAGE under reducing (R) condition. The gel was stained overnight with Coomassie Blue. The purity of the protein is greater than 90%.


Bioactivity-SPR**SEC-MALS**

The purity of Canine CD64, His Tag(Cat. No. FCA-C52H8) is more than 90% and the molecular weight of this protein is around 33-49 kDa verified by SEC-MALS.


[Report](#)



Canine CD64, His Tag (Cat. No. FCA-C52H8) captured on CM5 chip via anti-His antibody can bind Herceptin with an affinity constant of 2.38 nM as determined in a SPR assay (Biacore 8K) (Routinely tested).



Canine CD64, His Tag (Cat. No. FCA-C52H8) captured on CM5 chip via anti-His antibody can bind Ipilimumab with an affinity constant of 2.30 nM as determined in a SPR assay (Biacore 8K) (QC tested).



Canine CD64, His Tag (Cat. No. FCA-C52H8) captured on CM5 chip via anti-His antibody can bind Monoclonal Anti-Human CD3 Antibody, Mouse IgG2a (Clone: OKT3) with an affinity constant of 0.769 nM as determined in a SPR assay (Biacore 8K) (Routinely tested).

Background

Receptors that recognize the Fc portion of IgG are divided into three groups designated Fc gamma RI, RII, and RIII, also known respectively as CD64, CD32, and CD16. Fc gamma RI binds IgG with high affinity and functions during early immune responses. Fc gamma RII and RIII are low affinity receptors that recognize IgG as aggregates surrounding multivalent antigens during late immune responses. High affinity immunoglobulin gamma Fc receptor I is also known as FCGR1A, FCG1, FCGR1, CD64 and IGFR1, is a type of integral membrane glycoprotein that binds monomeric IgG-type antibodies with high affinity, which belongs to the immunoglobulin superfamily or FCGR1 family. FCGR1A / CD64 contains 3 Ig-like C2-type (immunoglobulin-like) domains. CD64 is constitutively found on only macrophages and monocytes, but treatment of polymorphonuclear leukocytes with cytokines like IFN γ and G-CSF can induce CD64 expression on these cells.

Clinical and Translational Updates

Please contact us via TechSupport@acrobiosystems.com if you have any question on this product.