

Mouse FOLR1 Protein, His Tag

Catalog # FO1-M5225



Synonym

FOLR-1,FBP,FOLR,FR α

Source

Mouse FOLR1, His Tag(FO1-M5225) is expressed from human 293 cells (HEK293). It contains AA Thr 25 - Ser 232 (Accession # [P35846-1](#)).
Predicted N-terminus: Thr 25

Molecular Characterization

FOLR1(Thr 25 - Ser 232)
P35846-1

Poly-his

This protein carries a polyhistidine tag at the C-terminus.
The protein has a calculated MW of 26.2 kDa. The protein migrates as 32-34 kDa and 35-40 kDa under reducing (R) condition (SDS-PAGE) due to glycosylation.

Endotoxin

Less than 1.0 EU per μ g by the LAL method / rFC method.

Purity

>95% as determined by SDS-PAGE.

Formulation

Lyophilized from 0.22 μ m filtered solution in Tris and Glycine, pH7.5 with trehalose as protectant.

Contact us for customized product form or formulation.

Reconstitution

Please see Certificate of Analysis for specific instructions.

For best performance, we strongly recommend you to follow the reconstitution protocol provided in the CoA.

Storage


For long term storage, the product should be stored at lyophilized state at -20°C or lower.

Please avoid repeated freeze-thaw cycles.

This product is stable after storage at:

- 20°C to -70°C for 12 months in lyophilized state;
- 70°C for 3 months under sterile conditions after reconstitution.

SDS-PAGE




Mouse FOLR1, His Tag on SDS-PAGE under reducing (R) condition. The gel was stained with Coomassie Blue. The purity of the protein is greater than 95%.

Bioactivity-ELISA



Mouse FOLR1 Protein, His Tag

Catalog # FO1-M5225



Immobilized Folic acid-BSA conjugate at 1 µg/mL (100 µL/well) can bind Mouse FOLR1, His Tag (Cat. No. FO1-M5225) with a linear range of 0.3-10 ng/mL (QC tested).

Background

Folate Receptor 1 (FOLR1) is also known as Folate receptor alpha, Folate Binding Protein (FBP), FOLR, and is a member of the folate receptor (FOLR) family. Members of this gene family have a high affinity for folic acid and for several reduced folic acid derivatives, and mediate delivery of 5-methyltetrahydrofolate to the interior of cells. Mature FOLR1 is an N-glycosylated protein that is anchored to the cell surface by a GPI linkage. FOLR1 is predominantly expressed on epithelial cells and is dramatically upregulated on many carcinomas. FOLR1 is internalized to the endosomal system where it dissociates from its ligand before recycling to the cell surface. A soluble form of FOLR1 can be proteolytically shed from the cell surface into the serum and breast milk. Defects in FOLR1 are the cause of neurodegeneration due to cerebral folate transport deficiency (NCFTD). NCFTD is an autosomal recessive disorder resulting from brain-specific folate deficiency early in life.

Discounts, Gifts,
and more!

