


**Synonym**

GPA33,A33

**Source**

Cynomolgus GPA33, His Tag(GP3-C52H3) is expressed from human 293 cells (HEK293). It contains AA Ile 22 - Val 235 (Accession # [G7NU40-1](#)).

Predicted N-terminus: Ile 22

**Molecular Characterization**

GPA33(Ile 22 - Val 235)	Poly-his
G7NU40-1	

This protein carries a polyhistidine tag at the C-terminus.

The protein has a calculated MW of 25.6 kDa. The protein migrates as 35-38 kDa and 40-45 kDa under reducing (R) condition (SDS-PAGE) due to glycosylation.

**Endotoxin**

Less than 1.0 EU per  $\mu$ g by the LAL method / rFC method.

**Purity**

>90% as determined by SDS-PAGE.

>95% as determined by SEC-MALS.

**Formulation**

Lyophilized from 0.22  $\mu$ m filtered solution in PBS, pH7.4 with trehalose as protectant.

Contact us for customized product form or formulation.

**Reconstitution**

Please see Certificate of Analysis for specific instructions.

*For best performance, we strongly recommend you to follow the reconstitution protocol provided in the CoA.*


**Storage**

For long term storage, the product should be stored at lyophilized state at -20°C or lower.


*Please avoid repeated freeze-thaw cycles.*

This product is stable after storage at:

- -20°C to -70°C for 12 months in lyophilized state;
- -70°C for 3 months under sterile conditions after reconstitution.

**SDS-PAGE**


Cynomolgus GPA33, His Tag on SDS-PAGE under reducing (R) condition. The gel was stained with Coomassie Blue. The purity of the protein is greater than 90%.

**SEC-MALS**


The purity of Cynomolgus GPA33, His Tag (Cat. No. GP3-C52H3) is more than 95% and the molecular weight of this protein is around 30-40 kDa verified by SEC-MALS.


[Report](#)

**Bioactivity-ELISA**

Discounts, Gifts,  
and more!



» [www.acrobiosystems.com](http://www.acrobiosystems.com)



Immobilized Cynomolgus GPA33, His Tag (Cat. No. GP3-C52H3) at 1 µg/mL (100 µL/well) can bind Monoclonal Anti-Human GPA33 Antibody, Human IgG1 with a linear range of 0.2-2 ng/mL (QC tested).

## Background

Glycoprotein A33 (GPA33) is also known as Cell surface A33 antigen, is a single-pass type I membrane protein which is expressed in normal gastrointestinal epithelium and in 95% of colon cancers. GPA33 The predicted mature protein has a 213-amino acid extracellular region, a single transmembrane domain, and a 62-amino acid intracellular tail. The sequence of the extracellular region contains 1 Ig-like C2-type (immunoglobulin-like) domain and 1 Ig-like V-type (immunoglobulin-like) domain characteristic of the CD2 subgroup of the immunoglobulin (Ig) superfamily, which contains. GPA33 may play a role in cell-cell recognition and signaling.

**Discounts, Gifts,  
and more!**



» [www.acrobiosystems.com](http://www.acrobiosystems.com)