



## Synonym

GPC3, OCI5, Glycan-3, GTR2-2, MXR7, DGSX, SDYS, SGB, SGBS, SGBS1

## Source

APC-Labeled Human Glycan 3 Protein, His Tag (GP3-HA2H7) is produced via conjugation of APC to Human Glycan 3 Protein, His Tag with a new generation site-specific technology under Star Staining labeling platform. Human Glycan 3 Protein, His Tag is expressed from human 293 cells (HEK293). It contains AA Gln 25 - His 559 (Accession # [P51654-1](#)).

Predicted N-terminus: Gln 25

## Molecular Characterization

This protein carries a polyhistidine tag at the C-terminus.

## Conjugate

APC

Excitation Wavelength: 640 nm

Emission Wavelength: 661 nm

## Purity

>90% as determined by SDS-PAGE.

## Formulation

Lyophilized from 0.22 µm filtered solution in PBS, 0.2% BSA, pH7.4 with trehalose as protectant.

Contact us for customized product form or formulation.

## Reconstitution

Please see Certificate of Analysis for specific instructions.

*For best performance, we strongly recommend you to follow the reconstitution protocol provided in the CoA.*

## Storage

For long term storage, the product should be stored at lyophilized state at -20°C or lower.

*Please protect from light and avoid repeated freeze-thaw cycles.*

This product is stable after storage at:


- -20°C to -70°C for 12 months in lyophilized state;
- -70°C for 3 months under sterile conditions after reconstitution.

**Star Staining** fluorescent-labeled products are developed by a new-generation site-specific labeling technology with Star Standard quality at ACROBiosystems

- ★ Using new-generation site-specific labeling technology to maintain natural bioactivity.
- ★ No non-specific binding to non-transduced PBMCs.
- ★ High specificity and sensitivity verified by flow cytometry.
- ★ High homogeneity and high batch-to-batch consistency.

## Evaluation of CAR expression

### FACS Analysis of Anti-Glycan 3 CAR Expression



Discounts, Gifts,  
and more!




» [www.acrobiosystems.com](http://www.acrobiosystems.com)



5e5 of anti-GPC3 CAR-293 cells were stained with 100  $\mu$ L of 1:50 dilution (2  $\mu$ L stock solution in 100  $\mu$ L FACS buffer) of APC-Labeled Human Glycan 3 Protein, His Tag (Cat. No. GP3-HA2H7) and negative control protein respectively (Fig. C and B), and non-transfected 293 cells were used as a control (Fig. A). APC signal was used to evaluate the binding activity (QC tested).

FACS Analysis of Non-specific binding to PBMCs



5e5 of PBMCs were stained with APC-Labeled Human Glycan 3 Protein, His Tag (Cat. No. GP3-HA2H7) and anti-CD3 antibody, washed and then analyzed with FACS. FITC signal was used to evaluate the expression of CD3+ T cells in PBMCs, and APC signal was used to evaluate the non-specific binding activity to PBMCs (QC tested).

## Background

Glycan-3 (GPC3) is also known as Intestinal protein OCI-5, GTR2-2, MXR7, which belongs to the glycan family. Glycan 3 / GPC-3 is highly expressed in lung, liver and kidney. Glycan-3 inhibits the dipeptidyl peptidase activity of DPP4. Glycan-3 may be involved in the suppression/modulation of growth in the predominantly mesodermal tissues and organs, and also may play a role in the modulation of IGF2 interactions with its receptor and thereby modulate its function.

Discounts, Gifts,  
and more!



» [www.acrobiosystems.com](http://www.acrobiosystems.com)