



Synonym

GPC3, OCI5, Glycan-3, GTR2-2, MXR7, DGSX, SDYS, SGB, SGBS, SGBS1

Source

Mouse Glycan 3, Fc Tag(GP3-M5254) is expressed from human 293 cells (HEK293). It contains AA Gln 25 - Arg 357 & Ser 358 - Met 557 (Accession # [Q8CFZ4-1](#)).

Predicted N-terminus: Gln 25 & Ser 358

Molecular Characterization

Glycan 3(Gln 25 - Met 557)	Fc(Pro 100 - Lys 330)
Q8CFZ4-1	P01857

This protein carries a human IgG1 Fc tag at the C-terminus.

The protein has a calculated MW of 38.0 kDa & 49.1 kDa. The protein migrates as 40 kDa, 60 kDa and 100 kDa under reducing (R) condition (SDS-PAGE) due to glycosylation.

Endotoxin

Less than 1.0 EU per μ g by the LAL method / rFC method.

Purity

>90% as determined by SDS-PAGE.

Formulation

Lyophilized from 0.22 μ m filtered solution in Tris with Glycine, Arginine and NaCl, pH7.5 with trehalose as protectant.

Contact us for customized product form or formulation.

Reconstitution

Please see Certificate of Analysis for specific instructions.

For best performance, we strongly recommend you to follow the reconstitution protocol provided in the CoA.

Storage


For long term storage, the product should be stored at lyophilized state at -20°C or lower.

Please avoid repeated freeze-thaw cycles.

This product is stable after storage at:

- -20°C to -70°C for 12 months in lyophilized state;
- -70°C for 3 months under sterile conditions after reconstitution.

SDS-PAGE



Mouse Glycan 3, Fc Tag on SDS-PAGE under reducing (R) condition. The gel was stained with Coomassie Blue. The purity of the protein is greater than 90%.

Bioactivity-ELISA

Discounts, Gifts,
and more!




» www.acrobiosystems.com

Mouse Glycan 3 / GPC3 Protein, Fc Tag

Catalog # GP3-M5254



BIOSYSTEMS
Acro
Surprise Inside!



Immobilized Human FGF basic, premium grade (Cat. No. BFF-H4117) at 2 μ g/mL (100 μ L/well) can bind Mouse Glycan 3, Fc Tag (Cat. No. GP3-M5254) with a linear range of 1-20 ng/mL (QC tested).

Background

Glycan-3 (GPC3) is also known as Intestinal protein OCI-5, GTR2-2, MXR7, which belongs to the glycan family. Glycan 3 / GPC-3 is highly expressed in lung, liver and kidney. Glycan-3 inhibits the dipeptidyl peptidase activity of DPP4. Glycan-3 may be involved in the suppression/modulation of growth in the predominantly mesodermal tissues and organs, and also may play a role in the modulation of IGF2 interactions with its receptor and thereby modulate its function.

Discounts, Gifts,
and more!



» www.acrobiosystems.com

4/21/2025