

**Synonym**

HGF, HPTA, SF

Source

Human HGF Protein, His Tag (HGF-H52H3) is expressed from human 293 cells (HEK293). It contains AA Gln 32 - Ser 728 (Accession # [P14210-1](#)).

Predicted N-terminus: Gln 32

Molecular Characterization

HGF(Gln 32 - Ser 728)
P14210-1 **Poly-his**

This protein carries a polyhistidine tag at the C-terminus.

The mature form of HGF is a disulfide-linked heterodimer composed of proteolytically cleaved α and β chain. The protein has a calculated MW of 81.6 kDa (α chain 53.7 kDa and β chain 27.9 kDa). The protein migrates as 80-90 kDa (alpha & Beta chain), 58-63 kDa (alpha chain), 33 kDa (Beta chain) when calibrated against [Star Ribbon Pre-stained Protein Marker](#) under reducing (R) condition (SDS-PAGE) due to glycosylation.

Endotoxin

Less than 0.1 EU per μ g by the LAL method / rFC method.

Sterility

Negative

Mycoplasma

Negative

Purity

>90% as determined by SDS-PAGE.

>95% as determined by SEC-MALS.

Formulation

Lyophilized from 0.22 μ m filtered solution in PBS, pH7.4 with trehalose as protectant.

Contact us for customized product form or formulation.

Reconstitution

Please see Certificate of Analysis for specific instructions.

For best performance, we strongly recommend you to follow the reconstitution protocol provided in the CoA.


Storage

For long term storage, the product should be stored at lyophilized state at -20°C or lower.


Please avoid repeated freeze-thaw cycles.

This product is stable after storage at:

- -20°C to -70°C for 12 months in lyophilized state;
- -70°C for 3 months under sterile conditions after reconstitution.

SDS-PAGE

Human HGF Protein, His Tag on SDS-PAGE under reducing (R) condition. The gel was stained with Coomassie Blue. The purity of the protein is greater than 90% (With [Star Ribbon Pre-stained Protein Marker](#)).

SEC-MALS


Discounts, Gifts,
 and more!




» www.acrobiosystems.com



The purity of Human HGF Protein, His Tag (Cat. No. HGF-H52H3) is more than 95% and the molecular weight of this protein is around 90-105 kDa verified by SEC-MALS.

[Report](#)**Bioactivity-Organoid Culture**

Human EGF (Cat. No. EGF-H52H3), Noggin (Cat. No. NON-H5257), R-spondin1 (Cat. No. RS6-H4220), FGF7 (Cat. No. FG7-H52H5), FGF10, HGF (Cat. No. HGF-H52H3) actively support liver ductal organoid growth.

Bioactivity-ELISA


Immobilized Human HGF Protein, His Tag (Cat. No. HGF-H52H3) at 2 µg/mL (100 µL/well) can bind Human HGF R, Fc Tag (Cat. No. MET-H5256) with a linear range of 1-39 ng/mL (QC tested).

Bioactivity-SPR


Discounts, Gifts,
and more!




» www.acrobiosystems.com



Human HGF R, Fc Tag (Cat. No. MET-H5256) captured on CM5 chip via Anti-human IgG Fc antibodies surface can bind Human HGF Protein, His Tag (Cat. No. HGF-H52H3) with an affinity constant of 0.486 nM as determined in a SPR assay (Biacore 8K) (Routinely tested).

Bioactivity-BLI

Loaded Human HGF R, Fc Tag (Cat. No. MET-H5256) on Protein A Biosensor, can bind Human HGF Protein, His Tag (Cat. No. HGF-H52H3) with an affinity constant of 3.29 nM as determined in BLI assay (ForteBio Octet Red96e) (Routinely tested).

Bioactivity-CELL BASE**Human HGF, His Tag stimulates the secretion of IL-11 by Saos-2 cells**

Human HGF Protein, His Tag (Cat. No. HGF-H52H3) stimulates the secretion IL-11 by Saos-2 cells. The specific activity of Human HGF Protein, His Tag is $>6.00 \times 10^5$ IU/mg, which is calibrated against WHO Hepatocyte Growth Factor (precursor) (Human rDNA derived) (NIBSC code: 96/556) (QC tested).

**Discounts, Gifts,
and more!****>> www.acrobiosystems.com**



Background

Hepatocyte growth factor (HGF) is a paracrine cellular growth, motility and morphogenic factor. Activating ligand for the receptor tyrosine kinase MET by binding to it and promoting its dimerization. Hepatocyte growth factor is secreted by mesenchymal cells and acts as a multi-functional cytokine on cells of mainly epithelial origin. Its ability to stimulate mitogenesis, cell motility, and matrix invasion gives it a central role in angiogenesis, tumorogenesis, and tissue regeneration. In addition, HGF has been implicated in a variety of cancers, including of the lungs, pancreas, thyroid, colon, and breast.

Discounts, Gifts,
and more!



» www.acrobiosystems.com