

Synonym


IgE

Source

Human IgE Fc, His Tag(IGE-H52H9) is expressed from human 293 cells (HEK293). It contains AA Ser 106 - Gly 427 (Accession # [P01854-1](#)).

Predicted N-terminus: Ser 106

Molecular Characterization



This protein carries a polyhistidine tag at the C-terminus.

The protein has a calculated MW of 37.6 kDa. The protein migrates as 45-50 kDa under reducing (R) condition (SDS-PAGE) due to glycosylation.

The protein is designed as a dimer.

Endotoxin

Less than 1.0 EU per µg by the LAL method / rFC method.

Purity

>95% as determined by SDS-PAGE.

>90% as determined by SEC-MALS.

Formulation

Lyophilized from 0.22 µm filtered solution in PBS, pH7.4 with trehalose as protectant.

Contact us for customized product form or formulation.

Reconstitution

Please see Certificate of Analysis for specific instructions.

For best performance, we strongly recommend you to follow the reconstitution protocol provided in the CoA.

Storage


For long term storage, the product should be stored at lyophilized state at -20°C or lower.

Please avoid repeated freeze-thaw cycles.

This product is stable after storage at:


- -20°C to -70°C for 12 months in lyophilized state;
- -70°C for 3 months under sterile conditions after reconstitution.

SDS-PAGE



Human IgE Fc, His Tag on SDS-PAGE under reducing (R) condition. The gel was stained with Coomassie Blue. The purity of the protein is greater than 95%.

SEC-MALS




The purity of Human IgE Fc, His Tag (Cat. No. IGE-H52H9) is more than 90% and the molecular weight of this protein is around 75-95 kDa verified by SEC-MALS.

[Report](#)


Bioactivity-ELISA






Immobilized Human Fc epsilon RI alpha, Fc Tag (Cat. No. FCA-H5259) at 5 µg/mL (100 µL/well) can bind Human IgE Fc, His Tag (Cat. No. IGE-H52H9) with a linear range of 0.6-10 ng/mL (QC tested).

Bioactivity-SPR



Human Fc epsilon RI alpha, Fc Tag (Cat. No. FCA-H5259) immobilized on CM5 Chip can bind Human IgE Fc, His Tag (Cat. No. IGE-H52H9) with an affinity constant of 0.585 nM as determined in a SPR assay (Biacore 8K) (Routinely tested).



Cynomolgus Fc epsilon RI alpha, His Tag (Cat. No. FCA-C52H9) immobilized on CM5 Chip can bind Human IgE Fc, His Tag (Cat. No. IGE-H52H9) with an affinity constant of 0.226 nM as determined in a SPR assay (Biacore 8K) (Routinely tested).

Background

As one of the five designated immunoglobulin isotypes, immunoglobulin E (IgE) plays a major role in atopic conditions by inducing immediate hypersensitivity reactions. IgE also contributes significantly to the body’s immune response to parasitic infections. IgE antibodies are predominantly found in the tissues, firmly attached to effector cells, such as mast cells and basophils, by high-affinity IgE Fc receptor (Fc epsilon RI) and low-affinity IgE receptor (Fc epsilon RII).

Discounts, Gifts,
and more!

