



Synonym


Integrin alpha V beta 1, ITGAV&ITGB1

Source

Cynomolgus Integrin alpha V beta 1 (ITGAV&ITGB1) Heterodimer Protein, His Tag&Tag Free (IT1-C52W4) is expressed from human 293 cells (HEK293). It contains AA Phe 31 - Pro 993 & Gln 161 - Asp 868 (Accession # [A0A2K5WCD3-1](#) & [A0A7N9D0D7](#)).

Predicted N-terminus: Phe 31 & Gln 161

Molecular Characterization



[Other Tags and Version](#) [Biotin & Other Labeled Version](#)

Cynomolgus Integrin alpha V beta 1 (ITGAV&ITGB1) Heterodimer Protein, His Tag&Tag Free, produced by co-expression of ITGAV and ITGB1, has a calculated MW of 113 kDa (ITGAV) and 83.6 kDa (ITGB1). Subunit ITGAV is fused with an acidic tail at the C-terminus and followed by a polyhistidine tag and subunit ITGB1 contains no tag but a basic tail at the C-terminus. The protein migrates as 110-130 kDa (ITGAV) and 80-100 kDa (ITGB1) when calibrated against [Star Ribbon Pre-stained Protein Marker](#) under non-reducing (NR) condition (SDS-PAGE) due to glycosylation.

Endotoxin

Less than 1.0 EU per μ g by the LAL method / rFC method.

Purity

>90% as determined by SDS-PAGE.

Formulation

Lyophilized from 0.22 μ m filtered solution in 50 mM Tris, 150 mM NaCl, pH8.0 with trehalose as protectant.

Contact us for customized product form or formulation.

Reconstitution

Please see Certificate of Analysis for specific instructions.

For best performance, we strongly recommend you to follow the reconstitution protocol provided in the CoA.

Storage

For long term storage, the product should be stored at lyophilized state at -20°C or lower.

Please avoid repeated freeze-thaw cycles.


This product is stable after storage at:

- -20°C to -70°C for 12 months in lyophilized state;
- -70°C for 3 months under sterile conditions after reconstitution.

ACRO Quality Management System

- [QMS\(ISO, GMP\)](#)
- [Quality Advantages](#)
- [Quality Control Process](#)

SDS-PAGE



Cynomolgus Integrin alpha V beta 1 (ITGAV&ITGB1) Heterodimer Protein, His Tag&Tag Free on SDS-PAGE under non-reducing (NR) condition. The gel was stained with Coomassie Blue. The purity of the protein is greater than 90% (With [Star Ribbon Pre-stained Protein Marker](#)).

Background

Integrin alpha-5/beta-1 is a receptor for fibrinogen. Integrin alpha-1/beta-1, alpha-2/beta-1, alpha-6/beta-1 and alpha-7/beta-1 are receptors for laminin. Integrin alpha-4/beta-1 is a receptor for VCAM1. It recognizes the sequence Q-I-D-S in VCAM1. Integrin alpha-9/beta-1 is a receptor for VCAM1, cytотactин and osteopontin. It recognizes the sequence A-E-I-D-G-I-E-L in cytотactин. Integrin alpha-V/beta-1 is also a receptor for vitronectin. Beta-1 integrins recognize the sequence R-G-D in a wide array of ligands. Isoform 2 interferes with isoform 1 resulting in a dominant negative effect on cell adhesion and migration (in vitro). When associated with alpha-7/beta-1 integrin, regulates cell adhesion and laminin matrix deposition.

Discounts, Gifts,
and more!



www.acrobiosystems.com

11/5/2025