

**Synonym**

L1R (MPXV)

Source

Monkeypox virus (strain Zaire-96-I-16) L1R, His Tag(L1R-M5241) is expressed from human 293 cells (HEK293). It contains AA Met 1 - Gln 152 (Accession # [Q8V4Z7](#)).

Predicted N-terminus: His

Molecular Characterization

Poly-his	L1R(Met 1 - Gln 152)
	Q8V4Z7

This protein carries a polyhistidine tag at the N-terminus.

The protein has a calculated MW of 19.7 kDa. The protein migrates as 18-19 kDa under reducing (R) condition (SDS-PAGE) due to glycosylation.

Endotoxin

Less than 1.0 EU per μ g by the LAL method / rFC method.

Purity

>95% as determined by SDS-PAGE.

>90% as determined by SEC-HPLC.

Formulation

Lyophilized from 0.22 μ m filtered solution in PBS, pH7.4 with trehalose as protectant.

Contact us for customized product form or formulation.

Reconstitution

Please see Certificate of Analysis for specific instructions.

For best performance, we strongly recommend you to follow the reconstitution protocol provided in the CoA.


Storage

For long term storage, the product should be stored at lyophilized state at -20°C or lower.


Please avoid repeated freeze-thaw cycles.

This product is stable after storage at:

- -20°C to -70°C for 12 months in lyophilized state;
- -70°C for 3 months under sterile conditions after reconstitution.

SDS-PAGE

Monkeypox virus (strain Zaire-96-I-16) L1R, His Tag on SDS-PAGE under reducing (R) condition. The gel was stained with Coomassie Blue. The purity of the protein is greater than 95%.

SEC-HPLC

The purity of Monkeypox virus (strain Zaire-96-I-16) L1R, His Tag (Cat. No. L1R-M5241) was greater than 90% as determined by SEC-HPLC.

Bioactivity-ELISA


Discounts, Gifts,
and more!



» www.acrobiosystems.com



Monkeypox virus (strain Zaire-96-I-16) L1R, His Tag ELISA
0.5 µg of Monkeypox virus (strain Zaire-96-I-16) L1R, His Tag per well



Immobilized Monkeypox virus (strain Zaire-96-I-16) L1R, His Tag (Cat. No. L1R-M5241) at 5 µg/mL (100 µL/well) can bind Anti-Monkeypox virus L1R Polyclonal Antibody, Rabbit, IgG with a linear range of 0.6-2 ng/mL (QC tested).

Background

Monkeypox is a rare zoonosis caused by monkeypox virus, which has become the most serious orthpoxvirus and consists of complex double stranded DNA. The cases are mostly in central and western Africa. The pathogenesis of monkeypox is that the virus invades the body from respiratory mucosa, multiplies in lymphocytes, and incurs into blood producing transient venereal toxemia. After the virus multiplies in cells, the cells can invade the blood and propagate to the skin of the whole body, causing lesions.

Discounts, Gifts,
and more!



» www.acrobiosystems.com