

Synonym

LAG3,CD223,FDC

Source

Canine LAG-3, His Tag(LA3-C52H7) is expressed from human 293 cells (HEK293). It contains AA Pro 23 - Leu 440 (Accession # [XP_038295087.1](#)).
Predicted N-terminus: Pro 23

Molecular Characterization

LAG-3(Pro 23 - Leu 440)
XP_038295087.1

Poly-his

This protein carries a polyhistidine tag at the C-terminus.

The protein has a calculated MW of 46.8 kDa. The protein migrates as 55-65 kDa under reducing (R) condition (SDS-PAGE) due to glycosylation.

The protein is designed as a dimer.

Endotoxin

Less than 1.0 EU per µg by the LAL method / rFC method.

Purity

>95% as determined by SDS-PAGE.

>90% as determined by SEC-MALS.

Formulation

Lyophilized from 0.22 µm filtered solution in PBS, pH7.4 with trehalose as protectant.

Contact us for customized product form or formulation.

Reconstitution

Please see Certificate of Analysis for specific instructions.

For best performance, we strongly recommend you to follow the reconstitution protocol provided in the CoA.

Storage


For long term storage, the product should be stored at lyophilized state at -20°C or lower.

Please avoid repeated freeze-thaw cycles.

This product is stable after storage at:


- 20°C to -70°C for 12 months in lyophilized state;
- 70°C for 3 months under sterile conditions after reconstitution.

SDS-PAGE



Canine LAG-3, His Tag on SDS-PAGE under reducing (R) condition. The gel was stained with Coomassie Blue. The purity of the protein is greater than 95%.

SEC-MALS



The purity of Canine LAG-3, His Tag (Cat. No. LA3-C52H7) is more than 90% and the molecular weight of this protein is around 96-116 kDa verified by SEC-MALS.

[Report](#)



Background

Lymphocyte activation gene 3 protein (LAG3) is also known as CD antigen CD223 and protein FDC, which belongs to immunoglobulin (Ig) superfamily and contains 4 extracellular Ig-like domains. The LAG3 gene contains 8 exons. The sequence data, exon/intron organization, and chromosomal localization all indicate a close relationship of LAG3 to CD4. LAG3 /CD223 involved in lymphocyte activation. LAG3 /CD223 binds to HLA class-II antigens.

Discounts, Gifts,
and more!

