

**Synonym**

LDLR, FH, FHC, LDLCQ2

Source

Rabbit LDL R, Fc Tag(LDR-R5255) is expressed from human 293 cells (HEK293). It contains AA Ala 9 - Gly 765 (Accession # [P20063-1](#)).

Molecular Characterization

LDL R(Ala 9 - Gly 765)	Fc(Pro 100 - Lys 330)
P20063-1	P01857

This protein carries a human IgG1 Fc tag at the C-terminus.

The protein has a calculated MW of 109.2 kDa. The protein migrates as 120-140 kDa under reducing (R) condition (SDS-PAGE) due to glycosylation.

Endotoxin

Less than 1.0 EU per μ g by the LAL method / rFC method.

Purity

>90% as determined by SDS-PAGE.

Formulation

Lyophilized from 0.22 μ m filtered solution in Tris with Glycine, Arginine and NaCl, pH7.5 with trehalose as protectant.

Contact us for customized product form or formulation.

Reconstitution

Please see Certificate of Analysis for specific instructions.

For best performance, we strongly recommend you to follow the reconstitution protocol provided in the CoA.


Storage

For long term storage, the product should be stored at lyophilized state at -20°C or lower.

Please avoid repeated freeze-thaw cycles.

This product is stable after storage at:

- -20°C to -70°C for 12 months in lyophilized state;
- -70°C for 3 months under sterile conditions after reconstitution.

SDS-PAGE


Rabbit LDL R, Fc Tag on SDS-PAGE under reducing (R) condition. The gel was stained with Coomassie Blue. The purity of the protein is greater than 90%.

Bioactivity-ELISA

Discounts, Gifts,
and more!




» www.acrobiosystems.com




Immobilized Human PCSK9 Propeptide & Rabbit Mature PCSK9 (154-686), His Tag at 2 µg/mL (100 µL/well) can bind Rabbit LDL R, Fc Tag (Cat. No. LDR-R5255) with a linear range of 2-78 ng/mL (QC tested).

Bioactivity-BLI



Loaded Rabbit LDL R, Fc Tag (Cat. No. LDR-R5255) on Protein A Biosensor, can bind Human Apolipoprotein E, His Tag (Cat. No. APE-H5246) with an affinity constant of 56.1 nM as determined in BLI assay (ForteBio Octet Red96e) (Routinely tested).



Loaded Rabbit LDL R, Fc Tag (Cat. No. LDR-R5255) on Protein A Biosensor, can bind Human Apolipoprotein E, His Tag (Cat. No. APE-5256) with an affinity constant of 38.5 nM as determined in BLI assay (ForteBio Octet Red96e) (Routinely tested).

Background

Low-Density Lipoprotein (LDL) Receptor is also known as LDLR, FH, FHC, LDLCQ2, and is a mosaic protein of ~840 amino acids (after removal of signal peptide) that mediates the endocytosis of cholesterol-rich LDL. It is a cell-surface receptor that recognizes the apoprotein B100 which is embedded in the phospholipid outer layer of LDL particles. The receptor also recognizes the apoE protein found in chylomicron remnants and VLDL remnants (IDL). It belongs to the Low density lipoprotein receptor gene family. LDL receptor complexes are present in clathrin-coated pits (or buds) on the cell surface, which when bound to LDL-cholesterol via adaptin, are pinched off to form clathrin-coated vesicles inside the cell. This allows LDL-cholesterol to be bound and internalized in a process known as endocytosis and prevents the LDL just diffusing around the membrane surface. This occurs in all nucleated cells (not erythrocytes), but mainly in the liver which removes ~70% of LDL from the circulation. Synthesis of receptors in the cell is regulated by the level of free intracellular cholesterol; if it is in excess for the needs of the cell then the transcription of the receptor gene will be inhibited. LDL receptors are translated by ribosomes on the endoplasmic reticulum and are modified by the Golgi apparatus before travelling in vesicles to the cell surface. LDL is directly involved in the development of atherosclerosis, due to accumulation of LDL-cholesterol in the blood. Atherosclerosis is the process responsible for the majority of cardiovascular diseases.

Discounts, Gifts,
and more!

