

**Synonym**

Leucine-rich repeat-containing protein 15,LRRC15,LIB,hLib

SourceMouse LRRC15, His Tag(LR5-M5225) is expressed from human 293 cells (HEK293). It contains AA Tyr 22 - Gly 536 (Accession # [Q80X72-1](#)).**Molecular Characterization**

LRRC15(Tyr 22 - Gly 536) Q80X72-1	Poly-his
--------------------------------------	----------

This protein carries a polyhistidine tag at the C-terminus.

The protein has a calculated MW of 59.5 kDa. The protein migrates as 65-90 kDa under reducing (R) condition (SDS-PAGE) due to glycosylation.

EndotoxinLess than 1.0 EU per μ g by the LAL method / rFC method.**Purity**

>95% as determined by SDS-PAGE.

>90% as determined by SEC-MALS.

FormulationLyophilized from 0.22 μ m filtered solution in PBS, pH7.4 with trehalose as protectant.

Contact us for customized product form or formulation.

Reconstitution

Please see Certificate of Analysis for specific instructions.


*For best performance, we strongly recommend you to follow the reconstitution protocol provided in the CoA.***Storage**

For long term storage, the product should be stored at lyophilized state at -20°C or lower.


Please avoid repeated freeze-thaw cycles.

This product is stable after storage at:

- -20°C to -70°C for 12 months in lyophilized state;
- -70°C for 3 months under sterile conditions after reconstitution.


SDS-PAGE

Mouse LRRC15, His Tag on SDS-PAGE under reducing (R) condition. The gel was stained with Coomassie Blue. The purity of the protein is greater than 95%.

SEC-MALS

The purity of Mouse LRRC15, His Tag (Cat. No. LR5-M5225) is more than 90% and the molecular weight of this protein is around 68-92 kDa verified by SEC-MALS.

[Report](#)**Bioactivity-ELISA**Discounts, Gifts,
and more!» www.acrobiosystems.com



Immobilized Mouse LRRC15, His Tag (Cat. No. LR5-M5225) at 10 μ g/mL (100 μ L/well) can bind Samrotamab with a linear range of 1-10 ng/mL (QC tested).

Background

LRRC15 (Leucine-rich repeat-containing protein 15) is also known as LIB and hLib. LRRC15 is highly expressed in a variety of solid tumors. LRRC15 was expressed on stromal fibroblasts in many solid tumors (e.g., breast, head and neck, lung, pancreatic) as well as directly on a subset of cancer cells of mesenchymal origin (e.g., sarcoma, melanoma, glioblastoma). LRRC15 expression was induced by TGF β on activated fibroblasts (α SMA+) and on mesenchymal stem cells. These collective findings suggested LRRC15 as a novel CAF and mesenchymal marker with utility as a therapeutic target for the treatment of cancers with LRRC15-positive stromal desmoplasia or cancers of mesenchymal origin. ABBV-085 is a monomethyl auristatin E (MMAE)-containing antibody–drug conjugate (ADC) directed against LRRC15, and it demonstrated robust preclinical efficacy against LRRC15 stromal-positive/cancer-negative, and LRRC15 cancer-positive models as a monotherapy, or in combination with standard-of-care therapies.

Discounts, Gifts,
and more!



» www.acrobiosystems.com