

**Synonym**

MIS RII,MRRII,AMHR2,AMHR,MISR2,AMH type II receptor

Source

Human MIS RII Protein, Mouse IgG2a Fc Tag(MII-H5253) is expressed from human 293 cells (HEK293). It contains AA Pro 18 - Ser 144 (Accession # [Q16671-1](#)).

Predicted N-terminus: Pro 18

Molecular Characterization

MIS RII(Pro 18 - Ser 144)	mFc(Glu 98 - Lys 330)
Q16671-1	P01863

This protein carries a mouse IgG2a Fc tag at the C-terminus. The protein has a calculated MW of 40.4 kDa. The protein migrates as 48-55 kDa when calibrated against [Star Ribbon Pre-stained Protein Marker](#) under reducing (R) condition (SDS-PAGE) due to glycosylation.

EndotoxinLess than 0.1 EU per μ g by the LAL method / rFC method.**Purity**

>90% as determined by SDS-PAGE.

FormulationLyophilized from 0.22 μ m filtered solution in PBS, pH7.4 with trehalose as protectant.

Contact us for customized product form or formulation.

Reconstitution

Please see Certificate of Analysis for specific instructions.


*For best performance, we strongly recommend you to follow the reconstitution protocol provided in the CoA.***Storage**

For long term storage, the product should be stored at lyophilized state at -20°C or lower.

Please avoid repeated freeze-thaw cycles.

This product is stable after storage at:

- -20°C to -70°C for 12 months in lyophilized state;
- -70°C for 3 months under sterile conditions after reconstitution.

SDS-PAGE

Human MIS RII Protein, Mouse IgG2a Fc Tag on SDS-PAGE under reducing (R) condition. The gel was stained with Coomassie Blue. The purity of the protein is greater than 90% (With [Star Ribbon Pre-stained Protein Marker](#)).

Background

MIS RII (Mullerian inhibiting substance type II receptor), also known as AMHRII (anti-Mullerian hormone type II receptor), is a serine/threonine receptor with a single transmembrane domain that belongs to the family of type II receptors of the TGF-beta superfamily. Mutations in MIS RII result in persistent Mullerian duct syndrome (PMDS), persistent Müllerian duct syndrome (PMDS) is a sex-limited disorder in which males develop portions of the female reproductive tract. Anti-

**Discounts, Gifts,
and more!****» www.acrobiosystems.com**

Human MIS RII Protein, Mouse IgG2a Fc Tag

Catalog # MII-H5253



BIOSYSTEMS
Acro
Surprise Inside!

Müllerian hormone (AMH) and its receptor (AMHR2) induce the regression of the Müllerian ducts in male embryos, but the mechanism by which the Amhr2 gene is specifically activated is not fully understood.

Discounts, Gifts,
and more!



» www.acrobiosystems.com

4/21/2025