



## Synonym

Mucin 1, MUC1, CD227, EMA, H23AG, KL-6, MAM6, MUC-1, SEC, MUC-1, X, MUC1, ZD, PEM, PEMT, PUM, CA15-3, Episialin

## Source

Human Mucin-1 (1098-1158), Mouse IgG2a Fc Tag (MU1-H5259) is expressed from human 293 cells (HEK293). It contains AA Ser 1098 - Gly 1158 (Accession # [P15941-1](#)).

Predicted N-terminus: Ser 1098

## Molecular Characterization

Mucin-1(Ser 1098 - Gly 1158)	mFc(Glu 98 - Lys 330)
P15941-1	P01863

This protein carries a mouse IgG2a Fc tag at the C-terminus.

The protein has a calculated MW of 33.4 kDa. The protein migrates as 43-53 kDa when calibrated against [Star Ribbon Pre-stained Protein Marker](#) under reducing (R) condition (SDS-PAGE) due to glycosylation.

## Endotoxin

Less than 0.1 EU per  $\mu$ g by the LAL method / rFC method.

## Purity

>95% as determined by SDS-PAGE.

>90% as determined by SEC-MALS.

## Formulation

Lyophilized from 0.22  $\mu$ m filtered solution in 50 mM Tris, 100 mM Glycine, 25 mM Arginine, 150 mM NaCl, pH7.5 with trehalose as protectant.

Contact us for customized product form or formulation.

## Reconstitution

Please see Certificate of Analysis for specific instructions.

*For best performance, we strongly recommend you to follow the reconstitution protocol provided in the CoA.*

## Storage


For long term storage, the product should be stored at lyophilized state at -20°C or lower.

*Please avoid repeated freeze-thaw cycles.*

This product is stable after storage at:


- 20°C to -70°C for 12 months in lyophilized state;
- 70°C for 3 months under sterile conditions after reconstitution.

## SDS-PAGE



Human Mucin-1 (1098-1158), Mouse IgG2a Fc Tag on SDS-PAGE under reducing (R) condition. The gel was stained with Coomassie Blue. The purity of the protein is greater than 95% (With [Star Ribbon Pre-stained Protein Marker](#)).

## SEC-MALS



The purity of Human Mucin-1 (1098-1158), Mouse IgG2a Fc Tag (Cat. No. MU1-H5259) is more than 90% and the molecular weight of this protein is around 68-88 kDa verified by SEC-MALS.

[Report](#)

Discounts, Gifts,  
and more!



» [www.acrobiosystems.com](http://www.acrobiosystems.com)



## Background

Membrane mucins have several functions in epithelial cells including cytoprotection, extravasation during metastases, maintenance of luminal structure, and signal transduction. MUC17, contains an extended, repetitive extracellular glycosylation domain and a carboxyl terminus with two EGF-like domains, a SEA module domain, a transmembrane domain, and a cytoplasmic domain with potential serine and tyrosine phosphorylation sites. Interacts via its C-terminus with PDZK1 and this interaction appears important for proper localization. Probably plays a role in maintaining homeostasis on mucosal surfaces.

Discounts, Gifts,  
and more!



» [www.acrobiosystems.com](http://www.acrobiosystems.com)