



Synonym

Neuraminidase/NA (Influenza Virus)

Source

Influenza A (A/Georgia/12/2022 CVR-167) Neuraminidase (NA) (A/pdm09 lineage) Protein, His Tag (NEE-V524p) is expressed from human 293 cells (HEK293). It contains AA Val 80- Asn 469 (Accession # EPI2756234, GISAID).

Predicted N-terminus: His

Molecular Characterization

Poly-his	Neuraminidase (NA)(Val 80- Asn 469)
	EPI2756234

This protein carries a polyhistidine tag at the N-terminus.

The protein has a calculated MW of 44.9 kDa. The protein migrates as 52-55 kDa when calibrated against [Star Ribbon Pre-stained Protein Marker](#) under reducing (R) condition (SDS-PAGE) due to glycosylation.

Endotoxin

Less than 1.0 EU per μ g by the LAL method / rFC method.

Purity

>90% as determined by SDS-PAGE.

Formulation

Lyophilized from 0.22 μ m filtered solution in PBS, pH7.4 with trehalose as protectant.

Contact us for customized product form or formulation.

Reconstitution

Please see Certificate of Analysis for specific instructions.

For best performance, we strongly recommend you to follow the reconstitution protocol provided in the CoA.

Storage


For long term storage, the product should be stored at lyophilized state at -20°C or lower.

Please avoid repeated freeze-thaw cycles.

This product is stable after storage at:

- -20°C to -70°C for 12 months in lyophilized state;
- -70°C for 3 months under sterile conditions after reconstitution.

SDS-PAGE



Influenza A (A/Georgia/12/2022 CVR-167) NA (A/pdm09 lineage) Protein, His Tag on SDS-PAGE under reducing (R) condition. The gel was stained with Coomassie Blue. The purity of the protein is greater than 90% (With [Star Ribbon Pre-stained Protein Marker](#)).

Bioactivity-ELISA


Discounts, Gifts,
and more!



» www.acrobiosystems.com



Influenza A (A/Georgia/12/2022 CVR-167) Neuraminidase (NA) (A/pdm09 lineage) Protein, His Tag ELISA
0.1 µg of Influenza A (A/Georgia/12/2022 CVR-167) Neuraminidase (NA) (A/pdm09 lineage) Protein, His Tag per well



Immobilized Influenza A (A/Georgia/12/2022 CVR-167) Neuraminidase (NA) (A/pdm09 lineage) Protein, His Tag (Cat. No. NEE-V524p) at 1 µg/mL (100 µL/well) can bind Anti-NA(A/PAN) Antibody, Mouse IgG with a linear range of 0.1-2 ng/mL (QC tested).

Background

Neuraminidase (NA) and hemagglutinin (HA) are major membrane glycoproteins found on the surface of influenza virus. Hemagglutinin binds to the sialic acid-containing receptors on the surface of host cells during initial infection and at the end of an infectious cycle. Neuraminidase, on the other hand, cleaves the HA-sialic acid bondage from the newly formed virions and the host cell receptors during budding. Neuraminidase thus is described as a receptor-destroying enzyme which facilitates virus release and efficient spread of the progeny virus from cell to cell.

Discounts, Gifts,
and more!



» www.acrobiosystems.com