

Source

Human metapneumovirus Post-fusion glycoprotein F0, His Tag(RSF-V52H5) is expressed from human 293 cells (HEK293). It contains AA Leu 19- Thr 489 (Accession # [Q6WB98](#)).

Molecular Characterization

Post-Fusion glycoprotein F0(Leu 19- Thr 489) Q6WB98	Poly-his
--	----------

This protein carries a polyhistidine tag at the C-terminus.

The protein has a calculated MW of 56.6 kDa. The protein migrates as 51-55 kDa when calibrated against [Star Ribbon Pre-stained Protein Marker](#) under reducing (R) condition (SDS-PAGE).

Endotoxin

Less than 1.0 EU per µg by the LAL method.

Purity

>90% as determined by SDS-PAGE.

>90% as determined by SEC-MALS.

Formulation

Lyophilized from 0.22 µm filtered solution in 0.1 M Sodium citrate, 0.5 M Arginine, pH5.5 with trehalose as protectant.

Contact us for customized product form or formulation.

Reconstitution

Please see Certificate of Analysis for specific instructions.

For best performance, we strongly recommend you to follow the reconstitution protocol provided in the CoA.

Storage


For long term storage, the product should be stored at lyophilized state at -20°C or lower.

Please avoid repeated freeze-thaw cycles.

This product is stable after storage at:

- 20°C to -70°C for 12 months in lyophilized state;
- 70°C for 3 months under sterile conditions after reconstitution.


SDS-PAGE



Human metapneumovirus Post-fusion glycoprotein F0, His Tag on SDS-PAGE under reducing (R) condition. The gel was stained with Coomassie Blue. The purity of the protein is greater than 90% (With [Star Ribbon Pre-stained Protein Marker](#)).


Bioactivity-ELISA

SEC-MALS




The purity of Human metapneumovirus Post-fusion glycoprotein F0, His Tag (Cat. No. RSF-V52H5) is more than 90% and the molecular weight of this protein is around 180-210 kDa verified by SEC-MALS. [Report](#)





Immobilized Human metapneumovirus Post-fusion glycoprotein F0, His Tag (Cat. No. RSF-V52H5) at 1 µg/mL (100 µL/well) can bind Anti-Fusion glycoprotein F0 Antibody, Mouse IgG2a (101F) with a linear range of 0.1-4 ng/mL and not bind Anti-F (RSV) MPE8, Human IgG1 (QC tested).

Bioactivity-SPR



Anti-Fusion glycoprotein F0 Antibody, Mouse IgG2a (101F) captured on Protein G-Series S sensor chip can bind Human metapneumovirus Post-fusion glycoprotein F0, His Tag (Cat. No. RSF-V52H5) with an affinity constant of 1.74 nM as determined in a SPR assay (Biacore 8K) (Routinely tested).

Background

Human respiratory syncytial virus (HRSV) is the most common etiological agent of acute lower respiratory tract disease in infants and can cause repeated infections throughout life. The RSV fusion glycoprotein (RSV F) is the principal target of RSV neutralizing antibodies in human sera. The RSV F is a type I viral fusion protein synthesized as inactive, single-chain polypeptides that assemble into trimers. RSV F fuses the viral and host cell membranes by irreversible protein refolding from the labile prefusion conformation to the stable post-fusion conformation. Both states exhibit epitopes targeted by neutralizing antibodies, and post-fusion RSV F is being developed as a vaccine candidate.

Clinical and Translational Updates

