

#### Limited Use & License Disclosure

# BY USE OF THIS PRODUCT, RESEARCHER AGREES TO BE BOUND BY THE FOLLOWING TERMS OF LIMITED USE OF THIS CELL LINE PRODUCT.

- If the researcher is not willing to accept the terms of limited use of this cell line product, and the product is unused, ACRO will accept return of the unused product.
- Researchers may use this product for research use only, no commercial use is allowed.

  "Commercial use" means any and all uses of this product and derivatives by a party for profit or other consideration and may include but is not limited to use in: (1) product manufacture; and (2) to provide a service, information or data; and/or resale of the product or its derivatives, whether or not such product or derivatives are resold for use in research.
- This cell line is neither intended for any animal or human therapeutic purposes nor for any direct human in vivo use. You have no right to share, modify, transfer, distribute, sell, sublicense, or otherwise make the cell line available for use to other researchers, laboratories, research institutions, hospitals, universities, or service organizations.
- ACROBIOSYSTEMS MAKES NO WARRANTIES OR REPRESENTATIONS OF ANY KIND, EITHER EXPRESSED OR IMPLIED, WITH RESPECT TO THE SUITABILITY OF THE CELL LINE FOR ANY PARTICULAR USE.
- ACROBIOSYSTEMS ACCEPTS NO LIABILITY IN CONNECTION WITH THE HANDLING OR USE OF THE CELL LINE.
- Modifications of the cell line, transfer to a third party, or commercial use of the cell line may
  require a separate license and additional fees. Please contact <u>order.cn@acrobiosystems.com</u> for
  further details.



# CHO/Human MRGPRX2 Stable Cell Line

| Catalog No.  | Size   |
|--------------|--|
| SCCHO-ATP215 | $2 \times (1 \text{ vial contains } \sim 5 \times 10^6 \text{ cells})$ |

## • Description

The CHO/Human MRGPRX2 Stable Cell Line was engineered to express the receptor full length human MRGPRX2 (Uniprot: Q96LB1). Surface expression of human MRGPRX2 was confirmed by flow cytometry.

# • Application

• Useful for cell-based MRGPRX2 binding assay

# • Cell Line Profile

| Cell line              | CHO/Human MRGPRX2 Stable Cell Line |  |
|------------------------|------------------------------------|--|
| Host Cell              | СНО                                |  |
| Property               | Adherent                           |  |
| Complete Growth Medium | F-12K + 10% FBS                    |  |
| Selection Marker       | Puromycin (2 μg/mL)                |  |
| Incubation             | 37°C with 5% CO <sub>2</sub>       |  |
| Doubling Time          | 22-24 hours                        |  |
| Transduction Technique | Lentivirus                         |  |



## • Materials Required for Cell Culture

• F-12K Nutrient Mixture (BasalMedia, Cat. No. L450KJ)

**Note:** If you are unable to obtain the specified F-12K Nutrient Mixture (BasalMedia, Cat. No. L450KJ) in China, you may use an alternative F-12K Nutrient Mixture (Gibco, Cat. No. 21127-022) or another suitable medium for culturing.

- Fetal bovine serum (CellMax, Cat. No. SA211.02)
- Puromycin (InvivoGen, Cat. No. ant-pr-5b)

**Note:** For selection antibiotics, we highly recommend using the specified brand. The activity of antibiotics may vary between manufacturers, so if you choose to use a different brand, it is essential to validate whether the concentration recommended in the culture medium is suitable. Regardless of the brand used, we recommend maintaining a backup culture without selection antibiotics to avoid potential cell loss due to inappropriate antibiotic concentration.

- 0.25% Trypsin-EDTA (1X), Phenol Red (Gibco, Cat. No. 25200-056)
- Penicillin-Streptomycin (Gibco, Cat. No. 15140-122)
- Phosphate Buffered Saline (1X) (HyClone, Cat. No. SH30256.01)
- Complete Growth Medium: F-12K + 10% FBS, 1%P/S
- Culture Medium: F-12K + 10% FBS, Puromycin (2 μg/mL), 1%P/S
- Freeze Medium: 90% FBS, 10% (V/V) DMSO
- T-75 Culture flask (Corning, Cat. No. 430641)
- Cryogenic storage vials (SARSTEDT, Cat. No. 72.379.007)
- Thermostat water bath
- Centrifuge (Cence, Model: L550)
- Cell counter (MONWEI, Model: SmartCell200A Plus)
- CO<sub>2</sub> Incubator (Thermo, Model: 3111)
- Biological Safety Cabinet (Thermo, Model: 1389)



#### • Recovery

- 1. Thaw the vial by gently agitating it in a 37°C water bath. To minimize the risk of contamination, ensure the cap remains out of the water. Thawing should be completed quickly, typically within 3-5 minutes.
- 2. After thawing, promptly remove the vial from the water bath and decontaminate it by spraying with 70% ethanol. From this point onward, all operations must be performed under strict aseptic conditions.
- 3. Transfer the contents of the vial to a centrifuge tube containing 4.0 mL of complete growth medium. Centrifuge at approximately 1000 rpm for 5 minutes.
- 4. Resuspend the cell pellet with 5 mL complete growth medium and transfer the cell suspension into a T-75 flask containing 10-15 mL of pre-warmed complete growth medium.
- 5. Incubate at 37°C with 5% CO<sub>2</sub> incubator until the cells are ready to be split.

#### • Subculture

- 1. Cell viability may be low after thawing, and full recovery may take up to a week. Monitor the cells daily until the culture reaches 80-90% confluency. At this point, remove and discard the spent medium. Avoid allowing the cells to become over-confluent to ensure optimal cell health.
- 2. Wash the cells once with sterile PBS. Avoid adding PBS directly onto the cell surface.
- 3. Add 3 mL of 0.25% Trypsin-EDTA to the T-75 flask. Place the flask at 37°C for 5-7 minutes, until 90% of the cells have detached. Monitor under a microscope to avoid over-trypsinization.
- 4. Add 6.0 to 8.0 mL of culture medium using a pipette and gently rinse the cells from the surface of the T-75 flask. Gently pipette up and down several times to achieve a single cell suspension without cell clumps.
- 5. Transfer appropriate aliquots of the cell suspension to a new T-75 flask. A subcultivation ratio of 1:6 to 1:10 is recommended. Adjust the ratio based on your specific culture system.
- 6. Incubate at 37°C with 5% CO<sub>2</sub> incubator.
- 7. When the cell culture reaches 80-90% confluency, proceed to the next subculture. Avoid over-confluency, as this may negatively impact cell performance in subsequent passages.

**Note:** After recovery, maintain the cells for 1-2 passages in the complete growth medium not containing the selection marker, if the cells are in good condition, transition to the culture medium containing the selection marker during subculturing.



#### • Cryopreservation

- 1. When the cell culture reaches 80-90% confluency, remove and discard the spent medium.
- 2. Wash the cells once with sterile PBS. Avoid adding PBS directly onto the cell surface.
- 3. Add 3 mL of 0.25% Trypsin-EDTA to the T-75 flask. Place the flask at 37°C for 5-7 minutes, until 90% of the cells have detached. Monitor under a microscope to avoid over-trypsinization.
- 4. Add 6.0 to 8.0 mL of complete growth medium using a pipette and gently rinse the cells from the surface of the T-75 flask. Gently pipette up and down several times to achieve a single cell suspension without cell clumps. Count the viable cells.
- 5. Transfer the cell suspension to a centrifuge tube. Centrifuge at 1000 rpm for 5 min at room temperature to pellet the cells.
- 6. After centrifugation, discard the supernatant. Resuspend the cells in ice cold freezing medium to a concentration of  $5\times10^6$  to  $1\times10^7$  cells/mL.
- 7. Aliquot the cell suspension into cryogenic storage vials. Place the vials in a programmable cooler or an insulated box placed in a -80°C freezer overnight, then transfer to liquid nitrogen storage for long-term storage.


Note: It is recommended to establish a cell bank at the earliest possible passage for long-term use.

### • Storage Condition

Cells must be received in a frozen state on dry ice and should be transferred to liquid nitrogen or a -80°C freezer immediately upon receipt. If stored in a -80°C freezer, it is recommended to limit the storage period to no more than two weeks. For long-term preservation, transfer the cells to liquid nitrogen is highly recommended.



## • Receptor Assay



| Catalog No.  | Stable Cell Line                   | MFI for MRGPRX2 (PE) |
|--------------|------------------------------------|----------------------|
| NA           | Negative Control Cell              | 58.86                |
| SCCHO-ATP215 | CHO/Human MRGPRX2 Stable Cell Line | 62574.83             |

**Fig1.** Expression analysis of human MRGPRX2 on CHO/Human MRGPRX2 Stable Cell Line by FACS. Cell surface staining was performed on CHO/Human MRGPRX2 Stable Cell Line or negative control cell using PE-labeled anti-human MRGPRX2 antibody.



# • Related Products

| <u>Products</u>   | Cat.No.      |
|---|--------------|
| CHO/Human LIGHT Stable Cell Line                              | SCCHO-ATP109 |
| CHO/Human BTLA Stable Cell Line                               | SCCHO-ATP110 |
| CHO/Human TSHR Stable Cell Line                               | SCCHO-ATP085 |
| CHO/Human LILRB4 Stable Cell Line                             | SCCHO-ATP087 |
| Raji/Membrane-Bound Human TL1A Stable Cell Line               | SCRAJ-STT204 |
| Human DR3 (TL1A receptor) (Luc) Jurkat Reporter Cell          | SCJUR-STF178 |
| Raji/Human HVEM Stable Cell Line                              | SCRAJ-STF108 |
| Human TSLP R (Luc) HEK293 Reporter Cell                       | CHEK-ATF045  |
| STAT3 (Luc) HEK293 Reporter Cell                              | CHEK-ATF047  |
| Human IL-5 R alpha/CD131 (Luc) HEK293 Reporter Cell           | CHEK-ATF074  |
| HEK293/Human OX40 / TNFRSF4 / CD134 Stable Cell Line          | CHEK-ATP053  |
| HEK293/Human OX40 Ligand / TNFSF4 Stable Cell Line            | CHEK-ATP054  |
| HEK293/Human FcRn (FCGRT & B2M) Stable Cell Line              | CHEK-ATP079  |
| Human IL-11 R alpha (Luc) HEK293 Reporter Cell                | CHEK-ATF052  |
| Human IL-4 R alpha/IL-13 R alpha 1 (Luc) HEK293 Reporter Cell | CHEK-ATF075  |
| Human IL-21 R/CD132 (Luc) HEK293 Reporter Cell                | CHEK-ATF051  |
| Human IL-31 RA/OSMR (Luc) HEK293 Reporter Cell                | CHEK-ATF094  |
| Human IL-10 R alpha/IL-10 R beta (Luc) HEK293 Reporter Cell   | CHEK-ATF095  |
| Human CD40 (Luc) HEK293 Reporter Cell                         | CHEK-ATF097  |
| Human IL-7 R alpha/CD132 (Luc) HEK293 Reporter Cell           | CHEK-ATF099  |
| NIH-3T3/Human IGF-1 R Stable Cell Line Development Service    | CNIH-ATP102  |
| Human HVEM (Luc) HEK293 Reporter Cell                         | CHEK-ATF105  |
| Human BTLA (Luc) Jurkat Reporter Cell                         | SCJUR-STF106 |
| Human IGF-1 R (Luc) HEK293 Reporter Cell                      | CHEK-ATF107  |
| Human GLP-2R (Luc) HEK293 Reporter Cell                       | CHEK-ATF128  |
| Human RANK (Luc) HEK293 Reporter Cell                         | CHEK-ATF129  |
| HEK293/FcRn (FCGRT & B2M), GFP Tag Stable Cell Line           | CHEK-ATP132  |
| HEK293/Human TSHR Stable Cell Line                            | CHEK-ATP086  |
|   |              |



## • Related Products

| <u>Products</u>   | Cat.No.     |
|---|-------------|
| HEK293/Human LILRB4 Stable Cell Line                          | CHEK-ATP088 |
| HEK293/Human TL1A Stable Cell Line                            | CHEK-ATP142 |
| Human IL-17 RA/IL-17 RC (Luc) HEK293 Reporter Cell            | CHEK-ATF133 |
| Human OX40 (Luc) HEK293 Reporter Cell                         | CHEK-ATF135 |
| Human IL-2 R beta/IL-2 R gamma (Luc) HEK293 Reporter Cell     | CHEK-ATF136 |
| HEK293/Human HVEM Stable Cell Line                            | CHEK-ATP147 |
| Human IL-23 R/IL-12 R beta 1(Luc) HEK293 Reporter Cell        | CHEK-ATF166 |
| Human IL-22 R alpha 1/IL-10 R beta (Luc) HEK293 Reporter Cell | CHEK-ATF167 |
| HEK293/Human CD40 Ligand / TNFSF5 Stable Cell Line            | CHEK-ATP041 |
| Human TSHR (Luc) HEK293 Reporter Cell                         | CHEK-ATF187 |
| Human PTH1R (Luc) HEK293 Reporter Cell                        | CHEK-ATF194 |
| HEK293/Membrane-Bound human TL1A Stable Cell Line             | CHEK-ATP198 |
| Human TACI (Luc) HEK293 Reporter Cell                         | CHEK-ATF197 |