

Human CD32b (Luc) Jurkat Reporter Cele Data Sheet

➤ *Limited Use & License Disclosure*

BY USE OF THIS PRODUCT, RESEARCHER AGREES TO BE BOUND BY THE FOLLOWING TERMS OF LIMITED USE OF THIS CELL LINE PRODUCT.

- If the researcher is not willing to accept the terms of limited use of this cell line product, and the product is unused, ACRO will accept return of the unused product.
- Researchers may use this product for research use only, no commercial use is allowed.
"Commercial use" means any and all uses of this product and derivatives by a party for profit or other consideration and may include but is not limited to use in: (1) product manufacture; and (2) to provide a service, information or data; and/or resale of the product or its derivatives, whether or not such product or derivatives are resold for use in research.
- This cell line is neither intended for any animal or human therapeutic purposes nor for any direct human in vivo use. You have no right to share, modify, transfer, distribute, sell, sublicense, or otherwise make the cell line available for use to other researchers, laboratories, research institutions, hospitals, universities, or service organizations.
- ACROBIOSYSTEMS MAKES NO WARRANTIES OR REPRESENTATIONS OF ANY KIND, EITHER EXPRESSED OR IMPLIED, WITH RESPECT TO THE SUITABILITY OF THE CELL LINE FOR ANY PARTICULAR USE.
- ACROBIOSYSTEMS ACCEPTS NO LIABILITY IN CONNECTION WITH THE HANDLING OR USE OF THE CELL LINE.
- Modifications of the cell line, transfer to a third party, or commercial use of the cell line may require a separate license and additional fees. Please contact order.cn@acrobiosystems.com for further details.

Human CD32b (Luc) Jurkat Reporter Cele Data Sheet

Human CD32b (Luc) Jurkat Reporter Cell


Catalog No.	Size
SCJUR-STF071	2 × (1 vial contains ~5×10 ⁶ cells)

• Description

The Human CD32b (Luc) Jurkat Reporter Cell was engineered to not only express the NFAT response element driving luciferase expressing systems, but also express the receptor full length human CD32b (Uniprot: P31994-1), which can use to evaluate ADCP activity of antibodies in the presence of corresponding target cells. When co-cultured with a target cell and relevant antibody, the antibody simultaneously binds the target cell antigen and CD32b receptor on the surface of Human CD32b (Luc) Jurkat Reporter Cell, resulting in receptor clustering, intracellular signaling and NFAT-mediated luminescence.

• Application

- Determination of ADCP activity induced by antibodies.



Human CD32b (Luc) Jurkat Reporter Cell Data Sheet

• Cell Line Profile

Cell line	Human CD32b (Luc) Jurkat Reporter Cell
Host Cell	Jurkat
Property	Suspension
Complete Growth Medium	RPMI-1640 + 10% FBS
Selection Marker	Puromycin (5 µg/mL) + Hygromycin B (20 µg/mL)
Incubation	37°C with 5% CO ₂
Doubling Time	16-20 hours
Transduction Technique	Lentivirus

• Materials Required for Cell Culture

- RPMI Medium 1640 (Gibco, Cat. No. 11875-093)
- Fetal bovine serum (CellMax, Cat. No. SA211.02)
- Puromycin (InvivoGen, Cat. No. ant-pr-5b)
- Hygromycin B (Invitrogen, Cat. No. 10687010)

Note: For selection antibiotics, we highly recommend using the specified brand. The activity of antibiotics may vary between manufacturers, so if you choose to use a different brand, it is essential to validate whether the concentration recommended in the culture medium is suitable. Regardless of the brand used, we recommend maintaining a backup culture without selection antibiotics to avoid potential cell loss due to inappropriate antibiotic concentration.

- Penicillin-Streptomycin (Gibco, Cat. No. 15140-122)
- Complete Growth Medium: RPMI-1640 + 10% FBS, 1%P/S
- Culture Medium: RPMI-1640 + 10% FBS, Puromycin (5 µg/mL), Hygromycin B (20 µg/mL), 1%P/S
- Freeze Medium: 90% FBS, 10% (V/V) DMSO
- T-75 Culture flask (Corning, Cat. No. 430641)
- Cryogenic storage vials (SARSTEDT, Cat. No. 72.379.007)
- Thermostat water bath
- Centrifuge (Cence, Model: L550)
- Cell counter (MONWEI, Model: SmartCell200A Plus)
- CO₂ Incubator (Thermo, Model: 3111)
- Biological Safety Cabinet (Thermo, Model: 1389)

Human CD32b (Luc) Jurkat Reporter Cell Data Sheet

• *Recovery*

1. Thaw the vial by gently agitating it in a 37°C water bath. To minimize the risk of contamination, ensure the cap remains out of the water. Thawing should be completed quickly, typically within 3-5 minutes.
2. After thawing, promptly remove the vial from the water bath and decontaminate it by spraying with 70% ethanol. From this point onward, all operations must be performed under strict aseptic conditions.
3. Transfer the contents of the vial to a centrifuge tube containing 4.0 mL of complete growth medium.
4. Count viable cells and centrifuge at approximately 1000 rpm for 5 minutes.
5. Discard the supernatant and resuspend the cell pellet in an appropriate amount of fresh **complete growth medium**. Adjust the cell density of the suspension to 1×10^6 viable cells/mL and transfer cells to an appropriate size vessel.
6. Incubate at 37°C with 5% CO₂ incubator.

• *Subculture*

Cell viability may be low after thawing, and full recovery (viability >90%) may take up to 1-2 weeks. Once the cell density reaches approximately 2×10^6 viable cells/mL, adjust the density to a range of 2×10^5 - 5×10^5 viable cells/mL by either adding the fresh **culture medium** or replacing the existing culture medium. Avoid allowing the cell density to exceed 3×10^6 cells/mL, as this may negatively impact cell performance in subsequent passages. T-75 flasks are recommended for subculturing.

• **Subculturing Frequency:** It is recommended to subculture every 3-4 days, adjusting the frequency based on the cell density in your specific culture system.

Note: After recovery, maintain the cells for 1-2 passages in the complete growth medium not containing the selection marker, if the cells are in good condition (viability >90%), transition to the culture medium containing the selection marker during subculturing.

Human CD32b (Luc) Jurkat Reporter Cell Data Sheet

• *Cryopreservation*

1. Count viable cells and harvest the cell suspension.
2. Centrifuge at 1000 rpm for 5 min at room temperature and resuspend cells in ice cold freezing medium to a concentration of 5×10^6 to 1×10^7 cells/mL.
3. Aliquot the cell suspension into cryogenic storage vials. Place the vials in a programmable cooler or an insulated box placed in a -80°C freezer overnight, then transfer to liquid nitrogen storage for long-term storage.

Note: It is recommended to establish a cell bank at the earliest possible passage for long-term use.

• *Storage*

Cells must be received in a frozen state on dry ice and should be transferred to liquid nitrogen or a -80°C freezer immediately upon receipt. If stored in a -80°C freezer, it is recommended to limit the storage period to no more than two weeks. For long-term preservation, transfer the cells to liquid nitrogen is highly recommended.

Human CD32b (Luc) Jurkat Reporter Cell Data Sheet

• Receptor Assay




Fig1. Expression analysis of human CD32b on Human CD32b (Luc) Jurkat Reporter Cell by FACS. Human CD32b (Luc) Jurkat Reporter Cell or negative control cell were stained with PE-Cy7-labeled anti-human CD32b antibody.

• Application

Determination of ADCP activity (RLU)




Fig2. ADCP response to anti-human CD20 antibody (RLU). Anti-human CD20 antibody-induced ADCP activity was evaluated using Human CD32b (Luc) Jurkat Reporter Cell in the presence of Raji cells that express CD20 endogenously. The EC₅₀ of anti-human CD20 antibody was approximately 0.112 $\mu\text{g/mL}$.

Human CD32b (Luc) Jurkat Reporter Cell Data Sheet





Fig3. ADCP response to anti-human CD20 antibody (FOLD). Anti-human CD20 antibody-induced ADCP activity was evaluated using Human CD32b (Luc) Jurkat Reporter Cell in the presence of Raji cells that express CD20 endogenously. The max induction fold was approximately 226.

• Passage Stability



Passage	MFI for CD32b (PE-Cy7)
P7	4051
P13	4371
P18	4189
P25	4465

Fig4. Passage stability analysis of receptors expression by FACS. Flow cytometry surface staining of human CD32b on Human CD32b (Luc) Jurkat Reporter Cell demonstrates consistent mean fluorescent intensity across passage 7-25.

Human CD32b (Luc) Jurkat Reporter Cell Data Sheet




Fig5. Passage stability analysis by Signaling Bioassay. The continuously growing Human CD32b (Luc) Jurkat Reporter Cell was stimulated with serial dilutions of anti-human CD20 antibody in the presence of Raji cells that express CD20 endogenously. Anti-human CD20 antibody stimulated response demonstrates passage stabilization (fold induction and EC50) across passage 7-25.

Human CD32b (Luc) Jurkat Reporter Cell Data Sheet

• *Related Products*

Products

Human CD16a (158V) (Luc) Jurkat Reporter Cell
 Human CD16a (158F) (Luc) Jurkat Reporter Cell
 Human CD32a (131H) (Luc) Jurkat Reporter Cell
 Human CD32a (131R) (Luc) Jurkat Reporter Cell
 Human CD64 (Luc) Jurkat Reporter Cell

Cat.No.

SCJUR-STF067
 SCJUR-STF068
 SCJUR-STF069
 SCJUR-STF070
 SCJUR-STF072