

Synonym

SHPS1,SIRPA,CD172A,BIT,MFR,MYD1,P84,PTPNS1

Source

Mouse SIRP alpha Protein, Mouse IgG2a Fc Tag(SIA-M5252) is expressed from human 293 cells (HEK293). It contains AA Lys 32 - Asn 373 (Accession # [BAA13520.1](#) (G365D)).  
Predicted N-terminus: Lys 32

Molecular Characterization



This protein carries a mouse IgG2a Fc tag at the C-terminus.  
The protein has a calculated MW of 65.0 kDa. The protein migrates as 90-130 kDa under reducing (R) condition (SDS-PAGE) due to glycosylation.

Endotoxin

Less than 0.01 EU per µg by the LAL method / rFC method.

Purity

>95% as determined by SDS-PAGE.  
>95% as determined by SEC-MALS.

Formulation

Lyophilized from 0.22 µm filtered solution in 50 mM Tris, 100 mM Glycine, pH7.5 with trehalose as protectant.  
Contact us for customized product form or formulation.

Reconstitution


Please see Certificate of Analysis for specific instructions.  
*For best performance, we strongly recommend you to follow the reconstitution protocol provided in the CoA.*

Storage

For long term storage, the product should be stored at lyophilized state at -20°C or lower.  
*Please avoid repeated freeze-thaw cycles.*  
This product is stable after storage at:


- 20°C to -70°C for 12 months in lyophilized state;
- 70°C for 3 months under sterile conditions after reconstitution.

SDS-PAGE



Mouse SIRP alpha Protein, Mouse IgG2a Fc Tag on SDS-PAGE under reducing (R) condition. The gel was stained with Coomassie Blue. The purity of the protein is greater than 95%.


SEC-MALS




The purity of Mouse SIRP alpha Protein, Mouse IgG2a Fc Tag (Cat. No. SIA-M5252) is more than 95% and the molecular weight of this protein is around 160-195 kDa verified by SEC-MALS.  
[Report](#)

Bioactivity-ELISA





Immobilized Mouse SIRP alpha Protein, Mouse IgG2a Fc Tag (Cat. No. SIA-M5252) at 2 µg/mL (100 µL/well) can bind Mouse CD47, Fc Tag (Cat. No. CD7-M5251) with a linear range of 8-63 ng/mL (QC tested).



Immobilized Human CD47 at 10 µg/mL (100 µL/well) can bind Mouse SIRP alpha Protein, Mouse IgG2a Fc Tag (Cat. No. SIA-M5252) with a linear range of 0.02-0.313 µg/mL (Routinely tested).

Background

Tyrosine-protein phosphatase non-receptor type substrate 1 (SHPS1) is also known as CD172 antigen-like family member A (CD172a), Macrophage fusion receptor, MyD-1 antigen, Signal-regulatory protein alpha (SIRPA or SIRP alpha) or p84, is a member of the SIRP family, and also belongs to the immunoglobulin superfamily. SIRP alpha is Ubiquitous and highly expressed in brain. SIRPA / CD172a is immunoglobulin-like cell surface receptor for CD47 and acts as docking protein and induces translocation of PTPN6, PTPN11 and other binding partners from the cytosol to the plasma membrane. SIRPA / SHPS-1 supports adhesion of cerebellar neurons, neurite outgrowth and glial cell attachment and may play a key role in intracellular signaling during synaptogenesis and in synaptic function By similarity. SIRPA / MyD1 involved in the negative regulation of receptor tyrosine kinase-coupled cellular responses induced by cell adhesion, growth factors or insulin and mediates negative regulation of phagocytosis, mast cell activation and dendritic cell activation. CD47 binding prevents maturation of immature dendritic cells and inhibits cytokine production by mature dendritic cells.

