



Synonym

SLAMF7,CD319,CS1,CRACC,19A,FOAP-12

Source

FITC-Labeled Human SLAMF7, His Tag (Cat. No. SL7-HF2H7) is expressed from human HEK293 cells. It contains AA Ser 23 - Met 226 (Accession # [AAH27867](#)). It is the FITC labeled form of Human SLAMF7, His Tag (Cat. No. SL7-H5225).

Predicted N-terminus: Ser 23

Molecular Characterization

SLAMF7(Ser 23 - Met 226)	Poly-his
AAH27867	

This protein carries a polyhistidine tag at the C-terminus.

The protein has a calculated MW of 23.2 kDa. The protein migrates as 33-45 kDa under reducing (R) condition (SDS-PAGE) due to glycosylation.

Conjugate

FITC

Excitation source: 488 nm spectral line, argon-ion laser

Excitation Wavelength: 488 nm

Emission Wavelength: 535 nm

Labeling

The primary amines in the side chains of lysine residues and the N-terminus of the protein are conjugated with FITC using standard chemical labeling method. The residual FITC is removed by molecular sieve treatment during purification process.

Protein Ratio

The FITC to protein molar ratio is 3-4.5.

Purity

>90% as determined by SDS-PAGE.

Formulation

Lyophilized from 0.22 µm filtered solution in PBS, pH7.4 with trehalose as protectant.

Contact us for customized product form or formulation.

Reconstitution

Please see Certificate of Analysis for specific instructions.

For best performance, we strongly recommend you to follow the reconstitution protocol provided in the CoA.

Storage


For long term storage, the product should be stored at lyophilized state at -20°C or lower.

Please protect from light and avoid repeated freeze-thaw cycles.

This product is stable after storage at:

- -20°C to -70°C for 12 months in lyophilized state;
- -70°C for 3 months under sterile conditions after reconstitution.

SDS-PAGE



Discounts, Gifts,
and more!




» www.acrobiosystems.com



FITC-Labeled Human SLAMF7, His Tag on SDS-PAGE under reducing (R) condition. The gel was stained with Coomassie Blue. The purity of the protein is greater than 90%.

Bioactivity-FACS



2e5 of Anti-SLAMF7 CAR-293 cells were stained with 100 μ L of 10 μ g/mL of FITC-Labeled Human SLAMF7, His Tag (Cat. No. SL7-HF2H7) and negative control protein respectively, FITC signal was used to evaluate the binding activity (QC tested)

Background

SLAM family member 7 (SLAMF7) is also known as CD2-like receptor-activating cytotoxic cells (CRACC), Membrane protein FOAP-12, CD antigen CD319, Novel Ly9, Protein 19A, which is a single-pass type I membrane protein and a member of the CD2 family of cell surface receptors. SLAMF7 is expressed in spleen, lymph node, peripheral blood leukocytes, bone marrow, small intestine, stomach, appendix, lung and trachea. Isoform 1 of SLAMF7 mediates NK cell activation through a SH2D1A-independent extracellular signal-regulated ERK-mediated pathway. May play a role in lymphocyte adhesion. Isoform 3 of SLAMF7 does not mediate any NK cell activation.

**Discounts, Gifts,
and more!**



» www.acrobiosystems.com