



Synonym

SLAMF7,CD319,CS1,CRACC,19A,FOAP-12

Source

Mouse SLAMF7 Protein, Mouse IgG2a Fc Tag(SL7-M5252) is expressed from human 293 cells (HEK293). It contains AA Ala 22 - Gly 224 (Accession # [Q8BHK6-1](#)).

Predicted N-terminus: Ala 22

Molecular Characterization

SLAMF7(Ala 22 - Gly 224)	mFc(Glu 98 - Lys 330)
Q8BHK6-1	P01863

This protein carries a mouse IgG2a Fc tag at the C-terminus.

The protein has a calculated MW of 49.4 kDa. The protein migrates as 55-70 kDa when calibrated against [Star Ribbon Pre-stained Protein Marker](#) under reducing (R) condition (SDS-PAGE) due to glycosylation.

Endotoxin

Less than 0.1 EU per μ g by the LAL method / rFC method.

Purity

>95% as determined by SDS-PAGE.

Formulation

Lyophilized from 0.22 μ m filtered solution in 50 mM Tris, 100 mM Glycine, pH7.5 with trehalose as protectant.

Contact us for customized product form or formulation.

Reconstitution

Please see Certificate of Analysis for specific instructions.

For best performance, we strongly recommend you to follow the reconstitution protocol provided in the CoA.

Storage


For long term storage, the product should be stored at lyophilized state at -20°C or lower.

Please avoid repeated freeze-thaw cycles.

This product is stable after storage at:

- -20°C to -70°C for 12 months in lyophilized state;
- -70°C for 3 months under sterile conditions after reconstitution.

SDS-PAGE



Mouse SLAMF7 Protein, Mouse IgG2a Fc Tag on SDS-PAGE under reducing (R) condition. The gel was stained with Coomassie Blue. The purity of the protein is greater than 95% (With [Star Ribbon Pre-stained Protein Marker](#)).

Background

SLAM family member 7 (SLAMF7) is also known as CD2-like receptor-activating cytotoxic cells (CRACC), Membrane protein FOAP-12, CD antigen CD319, Novel Ly9, Protein 19A, which is a single-pass type I membrane protein and a member of the CD2 family of cell surface receptors. SLAMF7 is expressed in spleen, lymph node, peripheral blood leukocytes, bone marrow, small intestine, stomach, appendix, lung and trachea. Isoform 1 of SLAMF7 mediates NK cell activation

Discounts, Gifts,
and more!



» www.acrobiosystems.com



through a SH2D1A-independent extracellular signal-regulated ERK-mediated pathway. May play a role in lymphocyte adhesion. Isoform 3 of SLAMF7 does not mediate any NK cell activation.

Discounts, Gifts,
and more!



» www.acrobiosystems.com