



Synonym

HAI, HAI1, MANSC2

Source

Biotinylated Human SPINT1 Protein, His,Avitag(SP1-H82E5) is expressed from human 293 cells (HEK293). It contains AA Pro 37 - Glu 449 (Accession # [O43278-2](#)).

Predicted N-terminus: Pro 37

Molecular Characterization

SPINT1(Pro 37 - Glu 449)
O43278-2

Poly-his

Avi

This protein carries a polyhistidine tag at the C-terminus, followed by an Avi tag (Avitag™).

The protein has a calculated MW of 49.7 kDa. The protein migrates as 53-63 kDa when calibrated against [Star Ribbon Pre-stained Protein Marker](#) under reducing (R) condition (SDS-PAGE) due to glycosylation.

Labeling

Biotinylation of this product is performed using Avitag™ technology. Briefly, the single lysine residue in the Avitag is enzymatically labeled with biotin.

Protein Ratio

Passed as determined by the HABA assay / binding ELISA.

Purity

>95% as determined by SDS-PAGE.

>90% as determined by SEC-MALS.

Formulation

Lyophilized from 0.22 µm filtered solution in 50 mM Tris, 150 mM NaCl, pH7.5 with trehalose as protectant.

Contact us for customized product form or formulation.

Reconstitution

Please see Certificate of Analysis for specific instructions.

For best performance, we strongly recommend you to follow the reconstitution protocol provided in the CoA.

Storage


For long term storage, the product should be stored at lyophilized state at -20°C or lower.

Please avoid repeated freeze-thaw cycles.

This product is stable after storage at:


- -20°C to -70°C for 12 months in lyophilized state;
- -70°C for 3 months under sterile conditions after reconstitution.

SDS-PAGE



Biotinylated Human SPINT1 Protein, His,Avitag on SDS-PAGE under reducing (R) condition. The gel was stained with Coomassie Blue. The purity of the protein is greater than 95% (With [Star Ribbon Pre-stained Protein Marker](#)).

SEC-MALS



The purity of Biotinylated Human SPINT1 Protein, His,Avitag (Cat. No. SP1-H82E5) is more than 90% and the molecular weight of this protein is around 45-70 kDa verified by SEC-MALS.

[Report](#)

Discounts, Gifts,
and more!



» www.acrobiosystems.com



Background

The protein encoded by this gene is a member of the Kunitz family of serine protease inhibitors. The protein is a potent inhibitor specific for HGF activator and is thought to be involved in the regulation of the proteolytic activation of HGF in injured tissues. Alternative splicing results in multiple variants encoding different isoforms. [provided by RefSeq, Jul 2008]

Discounts, Gifts,
and more!



» www.acrobiosystems.com