



## Synonym

Spike, S protein RBD, Spike glycoprotein Receptor-binding domain, S glycoprotein RBD, Spike protein RBD

## Source

SARS-CoV-2 Spike RBD Protein, His Tag (BQ.1/Omicron) (SPD-C5243) is expressed from human 293 cells (HEK293). It contains AA Arg 319 - Lys 537 (Accession # [QHD43416.1](#) (G339D, S371F, S373P, S375F, T376A, D405N, R408S, K417N, N440K, K444T, L452R, N460K, S477N, T478K, E484A, F486V, Q498R, N501Y, Y505H)). The spike mutations are identified on the SARS-CoV-2 Omicron variant (Pango lineage: BQ.1).

Predicted N-terminus: Arg 319

## Molecular Characterization

This protein carries a polyhistidine tag at the C-terminus.

The protein has a calculated MW of 26.6 kDa. The protein migrates as 33-40 kDa under reducing (R) condition (SDS-PAGE) due to glycosylation.

## Endotoxin

Less than 1.0 EU per  $\mu$ g by the LAL method / rFC method.

## Purity

>90% as determined by SDS-PAGE.

>90% as determined by SEC-MALS.

## Formulation

Lyophilized from 0.22  $\mu$ m filtered solution in PBS, pH7.4 with trehalose as protectant.

Contact us for customized product form or formulation.

## Reconstitution

Please see Certificate of Analysis for specific instructions.

*For best performance, we strongly recommend you to follow the reconstitution protocol provided in the CoA.*

## Storage


For long term storage, the product should be stored at lyophilized state at -20°C or lower.

*Please avoid repeated freeze-thaw cycles.*

This product is stable after storage at:


- -20°C to -70°C for 12 months in lyophilized state;
- -70°C for 3 months under sterile conditions after reconstitution.

## SDS-PAGE



SARS-CoV-2 Spike RBD Protein, His Tag (BQ.1/Omicron) on SDS-PAGE under reducing (R) condition. The gel was stained with Coomassie Blue. The purity of the protein is greater than 90%.

## SEC-MALS



The purity of SARS-CoV-2 Spike RBD Protein, His Tag (BQ.1/Omicron) (Cat. No. SPD-C5243) is more than 90% and the molecular weight of this protein is around 28-42 kDa verified by SEC-MALS.



[Report](#)

## Bioactivity-ELISA


Discounts, Gifts,  
and more!



» [www.acrobiosystems.com](http://www.acrobiosystems.com)




Immobilized Human ACE2, Fc Tag (Cat. No. AC2-H5257) at 5 µg/mL (100 µL/well) can bind SARS-CoV-2 Spike RBD Protein, His Tag (BQ.1/Omicron) (Cat. No. SPD-C5243) with a linear range of 2-31 ng/mL (QC tested).



Immobilized SARS-CoV-2 Spike RBD Protein, His Tag (BQ.1/Omicron) (Cat. No. SPD-C5243) at 1 µg/mL (100 µL/well) can bind Anti-SARS-CoV-2 Spike RBD Broadly Neutralizing Antibody, Human IgG1 (AM359b) (Cat. No. SPD-M265) with a linear range of 0.1-3 ng/mL (Routinely tested).

Immobilized SARS-CoV-2 Spike RBD Protein, His Tag (BQ.1/Omicron) (Cat. No. SPD-C5243) at 1 µg/mL (100 µL/well) can bind Human ACE2, Fc Tag (Cat. No. AC2-H5257) with a linear range of 0.1-3 ng/mL (Routinely tested).



Immobilized SARS-CoV-2 Spike RBD Protein, His Tag (BQ.1/Omicron) (Cat. No. SPD-C5243) at 1 µg/mL (100 µL/well) can bind Anti-SARS-CoV-2 Spike RBD Antibody, Chimeric mAb, Human IgG1 (AM130) (Cat. No. S1N-M13A1) with a linear range of 0.1-1 ng/mL (Routinely tested).

## Background

It's been reported that coronavirus can infect the human respiratory epithelial cells through interaction with the human ACE2 receptor. The spike protein is a large type I transmembrane protein containing two subunits, S1 and S2. S1 mainly contains a receptor binding domain (RBD), which is responsible for recognizing the cell surface receptor. S2 contains basic elements needed for the membrane fusion. The S protein plays key parts in the induction of neutralizing-antibody and T-cell responses, as well as protective immunity.

**Discounts, Gifts,  
and more!**

