

Synonym

Spike,S protein RBD,Spike glycoprotein Receptor-binding domain,S glycoprotein RBD,Spike protein RBD

Source

SARS-CoV-2 S protein RBD (K417T,E484K,N501Y), Fc Tag (SPD-C5258) is expressed from human 293 cells (HEK293). It contains AA Arg 319 - Lys 537 (Accession # [QHD43416.1](#) (K417T, E484K, N501Y)). The K417T, E484K, N501Y mutations were identified in the SARS-CoV-2 Gamma variant (Pango lineage: P.1; other names: 20J/501Y.V3).
Predicted N-terminus: Arg 319

Molecular Characterization



This protein carries a human IgG1 Fc tag at the C-terminus.
The protein has a calculated MW of 51.1 kDa. The protein migrates as 55-60 kDa under reducing (R) condition (SDS-PAGE) due to glycosylation.

Endotoxin

Less than 1.0 EU per µg by the LAL method / rFC method.

Purity

>95% as determined by SDS-PAGE.
>90% as determined by SEC-MALS.

Formulation

Lyophilized from 0.22 µm filtered solution in 50 mM Tris, 100 mM Glycine, 25 mM Arginine, 150 mM NaCl, pH7.5 with trehalose as protectant.
Contact us for customized product form or formulation.

Reconstitution


Please see Certificate of Analysis for specific instructions.
For best performance, we strongly recommend you to follow the reconstitution protocol provided in the CoA.

Storage

For long term storage, the product should be stored at lyophilized state at -20°C or lower.
Please avoid repeated freeze-thaw cycles.
This product is stable after storage at:

- 20°C to -70°C for 12 months in lyophilized state;
- 70°C for 3 months under sterile conditions after reconstitution.


SDS-PAGE



SARS-CoV-2 S protein RBD (K417T, E484K, N501Y), Fc Tag on SDS-PAGE under reducing (R) condition. The gel was stained with Coomassie Blue. The purity of the protein is greater than 95%.


Bioactivity-ELISA

SEC-MALS




The purity of SARS-CoV-2 S protein RBD (K417T, E484K, N501Y), Fc Tag (Cat. No. SPD-C5258) is more than 90% and the molecular weight of this protein is around 110-130 kDa verified by SEC-MALS.
[Report](#)






Immobilized SARS-CoV-2 S protein RBD (K417T, E484K, N501Y), Fc Tag (Cat. No. SPD-C5258) at 2 µg/mL (100 µL/well) can bind Human ACE2, His Tag (Cat. No. AC2-H52H8) with a linear range of 39-156 ng/mL (QC tested).



Immobilized Human ACE2, His Tag (Cat. No. AC2-H52H8) at 2 µg/mL (100 µL/well) can bind SARS-CoV-2 S protein RBD (K417T, E484K, N501Y), Fc Tag (Cat. No. SPD-C5258) with a linear range of 0.1-0.8 ng/mL (Routinely tested).

Bioactivity-BLI



Loaded SARS-CoV-2 S protein RBD, Fc Tag (Cat. No. SPD-C5258) on Protein A Biosensor, can bind Human ACE2, His Tag (Cat. No. AC2-H52H8) with an affinity constant of 2.05nM as determined in BLI assay (ForteBio Octet Red96e)(Routinely tested).

Background

It's been reported that Coronavirus can infect the human respiratory epithelial cells through interaction with the human ACE2 receptor. The spike protein is a large type I transmembrane protein containing two subunits, S1 and S2. S1 mainly contains a receptor binding domain (RBD), which is responsible for recognizing the cell surface receptor. S2 contains basic elements needed for the membrane fusion. The S protein plays key parts in the induction of neutralizing-antibody and T-cell responses, as well as protective immunity.

