



## Synonym

S1 protein NTD, Spike protein S1 NTD, BetaCoV S1-NTD

## Source

Biotinylated SARS-CoV-2 Spike NTD, His,Avitag (XBB.1/Omicron) (SPD-C82H7) is expressed from human 293 cells (HEK293). It contains AA Ser 13 - Leu 303 (Accession # [QHD43416.1](#) (T19I, LPP24-26del, A27S, V83A, G142D, Y144del, H146Q, Q183E, V213E, G252V)). The spike mutations are identified on the SARS-CoV-2 Omicron variant (Pango lineage: XBB.1).

Predicted N-terminus: Ser 13

## Molecular Characterization

This protein carries a polyhistidine tag at the C-terminus, followed by an Avi tag (Avitag™).

The protein has a calculated MW of 36.2 kDa. The protein migrates as 55-65 kDa under reducing (R) condition (SDS-PAGE) due to glycosylation.

## Labeling

*Biotinylation of this product is performed using Avitag™ technology. Briefly, the single lysine residue in the Avitag is enzymatically labeled with biotin.*

## Protein Ratio

Passed as determined by the HABA assay / binding ELISA.

## Purity

>90% as determined by SDS-PAGE.

>95% as determined by SEC-MALS.

## Formulation

Lyophilized from 0.22 µm filtered solution in PBS, pH7.4 with trehalose as protectant.

Contact us for customized product form or formulation.

## Reconstitution

Please see Certificate of Analysis for specific instructions.

*For best performance, we strongly recommend you to follow the reconstitution protocol provided in the CoA.*

## Storage


For long term storage, the product should be stored at lyophilized state at -20°C or lower.

*Please avoid repeated freeze-thaw cycles.*

This product is stable after storage at:


- -20°C to -70°C for 12 months in lyophilized state;
- -70°C for 3 months under sterile conditions after reconstitution.

## SDS-PAGE



Biotinylated SARS-CoV-2 Spike NTD, His,Avitag (XBB.1/Omicron) on SDS-PAGE under reducing (R) condition. The gel was stained with Coomassie Blue. The purity of the protein is greater than 90%.

## SEC-MALS



The purity of Biotinylated SARS-CoV-2 Spike NTD, His,Avitag (XBB.1/Omicron) (Cat. No. SPD-C82H7) is more than 95% and the molecular weight of this protein is around 50-60 kDa verified by SEC-MALS. [Report](#)

## Bioactivity-ELISA


Discounts, Gifts,  
and more!



» [www.acrobiosystems.com](http://www.acrobiosystems.com)



Biotinylated SARS-CoV-2 Spike NTD, His,Avitag (XBB.1/Omicron) ELISA  
 0.1  $\mu$ g of Biotinylated SARS-CoV-2 Spike NTD, His,Avitag (XBB.1/Omicron) per well



Immobilized Biotinylated SARS-CoV-2 Spike NTD, His,Avitag (XBB.1/Omicron) (Cat. No. SPD-C82H7) at 1  $\mu$ g/mL (100  $\mu$ L/well) on streptavidin (Cat. No. STN-N5116) precoated (0.5  $\mu$ g/well) plate can bind Anti-SARS-CoV-2 Spike NTD Antibody, Chimeric mAb, Human IgG1 (AM121) (Cat. No. SPD-M121) with a linear range of 0.1-2 ng/mL (QC tested).

## Background

It's been reported that coronavirus can infect the human respiratory epithelial cells through interaction with the human ACE2 receptor. The spike protein is a large type I transmembrane protein containing two subunits, S1 and S2. S1 mainly contains a receptor binding domain (RBD), which is responsible for recognizing the cell surface receptor. S2 contains basic elements needed for the membrane fusion. The S protein plays key parts in the induction of neutralizing-antibody and T-cell responses, as well as protective immunity.

Discounts, Gifts,  
 and more!



» [www.acrobiosystems.com](http://www.acrobiosystems.com)