

**Synonym**

Spike, S protein, Spike glycoprotein, S glycoprotein

Source


SARS-CoV-2 Spike Trimer, His Tag (CH.1.1/Omicron) (SPN-C5249) is expressed from human 293 cells (HEK293). It contains AA Val 16 - Pro 1213 (Accession # [QHD43416.1](#) (T19I, LPP24-26del, A27S, G142D, K147E, W152R, F157L, I210V, V213G, G257S, G339H, R346T, S371F, S373P, S375F, T376A, D405N, R408S, K417N, N440K, K444T, G446S, L452R, N460K, S477N, T478K, E484A, F486S, Q498R, N501Y, Y505H, D614G, H655Y, N679K, P681H, N764K, D796Y, Q954H, N969K, R683A, R685A, F817P, A892P, A899P, A942P, K986P, V987P)). The spike mutations are identified on the SARS-CoV-2 Omicron variant (Pango lineage: CH.1.1). The recombinant protein is expressed from human 293 cells (HEK293) with T4 fibritin trimerization motif and a polyhistidine tag at the C-terminus. Proline substitutions (F817P, A892P, A899P, A942P, K986P, V987P) and alanine substitutions (R683A and R685A) are introduced to stabilize the trimeric prefusion state of SARS-CoV-2 S protein and abolish the furin cleavage site, respectively.

Predicted N-terminus: Val 16

Molecular Characterization

This protein carries a polyhistidine tag at the C-terminus.

The protein has a calculated MW of 137.9 kDa. The protein migrates as 170-200 kDa under reducing (R) condition (SDS-PAGE) due to glycosylation.

EndotoxinLess than 1.0 EU per μ g by the LAL method / rFC method.**SDS-PAGE**

SARS-CoV-2 Spike Trimer, His Tag (CH.1.1/Omicron) on SDS-PAGE under reducing (R) condition. The gel was stained with Coomassie Blue. The purity of the protein is greater than 95%.

Purity

>95% as determined by SDS-PAGE.

>90% as determined by SEC-MALS.

FormulationLyophilized from 0.22 μ m filtered solution in 0.1 M Sodium citrate, pH5.5 with trehalose as protectant.

Contact us for customized product form or formulation.

Reconstitution

Please see Certificate of Analysis for specific instructions.


*For best performance, we strongly recommend you to follow the reconstitution protocol provided in the CoA.***Storage**

For long term storage, the product should be stored at lyophilized state at -20°C or lower.

Please avoid repeated freeze-thaw cycles.

This product is stable after storage at:

- -20°C to -70°C for 12 months in lyophilized state;
- -70°C for 3 months under sterile conditions after reconstitution.



SEC-MALS

The purity of SARS-CoV-2 Spike Trimer, His Tag (CH.1.1/Omicron) (Cat. No. SPN-C5249) is more than 90% and the molecular weight of this protein is around 470-520 kDa verified by SEC-MALS.

**Discounts, Gifts,
and more!**» www.acrobiosystems.com



[Report](#)

Bioactivity-ELISA



Immobilized Human ACE2, Fc Tag (Cat. No. AC2-H5257) at 5 µg/mL (100 µL/well) can bind SARS-CoV-2 Spike Trimer, His Tag (CH.1.1/Omicron) (Cat. No. SPN-C5249) with a linear range of 0.1-2 ng/mL (QC tested).

Immobilized SARS-CoV-2 Spike Trimer, His Tag (CH.1.1/Omicron) (Cat. No. SPN-C5249) at 1 µg/mL (100 µL/well) can bind Anti-SARS-CoV-2 Spike RBD Antibody, Chimeric mAb, Human IgG1 (AM130) (Cat. No. S1N-M13A1) with a linear range of 0.1-5 ng/mL (Routinely tested).



Immobilized SARS-CoV-2 Spike Trimer, His Tag (CH.1.1/Omicron) (Cat. No. SPN-C5249) at 1 µg/mL (100 µL/well) can bind Anti-SARS-CoV-2 Spike RBD Broadly Neutralizing Antibody, Human IgG1 (Cat. No. SPD-M265) with a linear range of 0.1-3 ng/mL (Routinely tested).

Background

It's been reported that coronavirus can infect the human respiratory epithelial cells through interaction with the human ACE2 receptor. The spike protein is a large type I transmembrane protein containing two subunits, S1 and S2. S1 mainly contains a receptor binding domain (RBD), which is responsible for recognizing the cell surface receptor. S2 contains basic elements needed for the membrane fusion. The S protein plays key parts in the induction of neutralizing antibody and T-cell responses, as well as protective immunity.

Discounts, Gifts,
and more!



» www.acrobiosystems.com