

**Synonym**

DDPAC,FTDP-17,MAPT,MSTD,MTBT1,Tau,PHF-tau,TAU

Source

Human Tau-441, His Tag(TAU-H51H3) is expressed from *E. coli* cells. It contains AA Met 1 - Leu 441 (Accession # [P10636-8](#)).

Predicted N-terminus: Met

Molecular Characterization

Poly-his

Tau-441(Met 1 - Leu 441)
P10636-8

This protein carries a polyhistidine tag at the N-terminus.

The protein has a calculated MW of 47.9 kDa. The protein migrates as 60 kDa under reducing (R) condition (SDS-PAGE).

Purity

>90% as determined by SDS-PAGE.

Formulation

Lyophilized from 0.22 μ m filtered solution in 50 mM Tris, 150 mM NaCl, pH7.5 with trehalose as protectant.

Contact us for customized product form or formulation.

Reconstitution

Please see Certificate of Analysis for specific instructions.

For best performance, we strongly recommend you to follow the reconstitution protocol provided in the CoA.


Storage

For long term storage, the product should be stored at lyophilized state at -20°C or lower.

Please avoid repeated freeze-thaw cycles.

This product is stable after storage at:

- -20°C to -70°C for 12 months in lyophilized state;
- -70°C for 3 months under sterile conditions after reconstitution.

SDS-PAGE


Human Tau-441, His Tag on SDS-PAGE under reducing (R) condition. The gel was stained with Coomassie Blue. The purity of the protein is greater than 90%.

Bioactivity-ELISA

Discounts, Gifts,
and more!



» www.acrobiosystems.com



Immobilized Human Tau-441, His Tag (Cat. No. TAU-H51H3) at 5 µg/mL (100 µL/well) can bind Tau Monoclonal Antibody with a linear range of 1-20 ng/mL (Routinely tested).

Background

Tau is a microtubule-associated protein, which encodes by the MAPT gene that located on chromosome 17q21. Tau Promotes microtubule assembly and stability, and might be involved in the establishment and maintenance of neuronal polarity. Hyperphosphorylation of the tau protein (tau inclusions, pTau) can result in the self-assembly of tangles of paired helical filaments and straight filaments, which are involved in the pathogenesis of Alzheimer's disease, frontotemporal dementia, and other tauopathies. Tau-441 is known as "2N4R," "Isoform Tau-F," "Tau-4" or "Tau 441", which consisting of 441 amino acid. Tau-441 is a potential therapeutic target for pathogenesis.

**Discounts, Gifts,
and more!**



» www.acrobiosystems.com