

**Synonym**

BGCAN, betaglycan

Source

Human TGF-beta RIII / TGFBR3 Protein, His Tag(TG3-H52H3) is expressed from human 293 cells (HEK293). It contains AA Gly 21 - Val 787 (Accession # [Q03167-1](#)).

Predicted N-terminus: Gly 21

Molecular Characterization

TGF-beta RIII(Gly 21 - Val 787)	Poly-his
Q03167-1	

This protein carries a polyhistidine tag at the C-terminus.

The protein has a calculated MW of 86.7 kDa. The protein migrates as 100 kDa and 130-170 kDa when calibrated against [Star Ribbon Pre-stained Protein Marker](#) under reducing (R) condition (SDS-PAGE) due to glycosylation.

Endotoxin

Less than 1.0 EU per μ g by the LAL method / rFC method.

Purity

>90% as determined by SDS-PAGE.

Formulation

Lyophilized from 0.22 μ m filtered solution in PBS, pH7.4 with trehalose as protectant.

Contact us for customized product form or formulation.

Reconstitution

Please see Certificate of Analysis for specific instructions.

For best performance, we strongly recommend you to follow the reconstitution protocol provided in the CoA.


Storage

For long term storage, the product should be stored at lyophilized state at -20°C or lower.

Please avoid repeated freeze-thaw cycles.


This product is stable after storage at:

- -20°C to -70°C for 12 months in lyophilized state;
- -70°C for 3 months under sterile conditions after reconstitution.

SDS-PAGE

Human TGF-beta RIII / TGFBR3 Protein, His Tag on SDS-PAGE under reducing (R) condition. The gel was stained with Coomassie Blue. The purity of the protein is greater than 90% (With [Star Ribbon Pre-stained Protein Marker](#)).

Bioactivity-BLI**Discounts, Gifts,
and more!****» www.acrobiosystems.com**



Loaded Biotinylated Human TGF-Beta 1, Avitag (Cat. No. TG1-H8217) on SA Biosensor, can bind Human TGF-beta RIII / TGFBR3 Protein, His Tag (Cat. No. TG3-H52H3) with an affinity constant of 1.80 nM as determined in BLI assay (ForteBio Octet Red96e) (QC tested).

Background

This locus encodes the transforming growth factor (TGF)-beta type III receptor. The encoded receptor is a membrane proteoglycan that often functions as a co-receptor with other TGF-beta receptor superfamily members. Ectodomain shedding produces soluble TGFBR3, which may inhibit TGF β signaling. Decreased expression of this receptor has been observed in various cancers. Alternatively spliced transcript variants encoding different isoforms have been identified for this gene.

**Discounts, Gifts,
and more!**



» www.acrobiosystems.com