



Synonym

TIGIT,VSIG9,VSTM3

Source

Biotinylated Cynomolgus / Rhesus macaque TIGIT Protein, Fc,Avitag(TIT-C82F3) is expressed from human 293 cells (HEK293). It contains AA Met 89 - Ile 208 (Accession # [G7NXM4-1](#)).
Predicted N-terminus: Met 89

Molecular Characterization

TIGIT(Met 89 - Ile 208) G7NXM4-1	Fc(Pro 100 - Lys 330) P01857	Avi
-------------------------------------	---------------------------------	-----

This protein carries a human IgG1 Fc tag at the C-terminus, followed by an Avi tag (Avitag™).
The protein has a calculated MW of 41.4 kDa. The protein migrates as 48-53 kDa when calibrated against [Star Ribbon Pre-stained Protein Marker](#) under reducing (R) condition (SDS-PAGE) due to glycosylation.

Labeling

Biotinylation of this product is performed using Avitag™ technology. Briefly, the single lysine residue in the Avitag is enzymatically labeled with biotin.

Protein Ratio

Passed as determined by the HABA assay / binding ELISA.

Purity

>90% as determined by SDS-PAGE.
>95% as determined by SEC-MALS.

Formulation

Lyophilized from 0.22 µm filtered solution in PBS, pH7.4 with trehalose as protectant.
Contact us for customized product form or formulation.

Reconstitution


Please see Certificate of Analysis for specific instructions.
For best performance, we strongly recommend you to follow the reconstitution protocol provided in the CoA.

Storage

For long term storage, the product should be stored at lyophilized state at -20°C or lower.
Please avoid repeated freeze-thaw cycles.
This product is stable after storage at:


- 20°C to -70°C for 12 months in lyophilized state;
- 70°C for 3 months under sterile conditions after reconstitution.

SDS-PAGE



Biotinylated Cynomolgus / Rhesus macaque TIGIT Protein, Fc,Avitag on SDS-PAGE under reducing (R) condition. The gel was stained with Coomassie Blue. The purity of the protein is greater than 90% (With [Star Ribbon Pre-stained Protein Marker](#)).

SEC-MALS




The purity of Biotinylated Cynomolgus / Rhesus macaque TIGIT Protein, Fc,Avitag (Cat. No. TIT-C82F3) is more than 95% and the molecular weight of this protein is around 90-110 kDa verified by SEC-MALS.
[Report](#)

Bioactivity-ELISA



Biotinylated Cynomolgus / Rhesus macaque TIGIT Protein, Fc,Avitag ELISA
0.1 µg of Biotinylated Cynomolgus / Rhesus macaque TIGIT Protein, Fc,Avitag per well



Immobilized Biotinylated Cynomolgus / Rhesus macaque TIGIT Protein, Fc,Avitag (Cat. No. TIT-C82F3) at 1 µg/mL (100 µL/well)on streptavidin precoated (STN-N5116) (can bind Mouse CD155, Mouse IgG2a Fc Tag (Cat. No. CD5-M5254) with a linear range of 0.004-0.125 µg/mL (QC tested).

Background

T-cell immunoreceptor with Ig and ITIM domains (TIGIT) is also known as V-set and immunoglobulin domain-containing protein 9 (VSIG9), V-set and transmembrane domain-containing protein 3 (VSTM3),which belongs to single-pass type I membrane protein containing an immunoglobulin variable domain, a transmembrane domain and an immunoreceptor tyrosine-based inhibitory motif (ITIM). TIGIT is expressed at low levels on peripheral memory and regulatory CD4+ T-cells and NK cells and is up-regulated following activation of these cells (at protein level). TIGIT binds with high affinity to the poliovirus receptor (PVR) which causes increased secretion of IL10 and decreased secretion of IL12B and suppresses T-cell activation by promoting the generation of mature immunoregulatory dendritic cells.

