

PE-Labeled Human TIGIT Protein, His TagStar Staining

Catalog # TIT-HP2H3



Synonym

TIGIT,VSIG9,VSTM3

Source

PE-Labeled Human TIGIT Protein, His Tag(TIT-HP2H3) is expressed from human 293 cells (HEK293). It contains AA Met 22 - Pro 141 (Accession # [Q495A1-1](#)).

Predicted N-terminus: Met 22

Molecular Characterization

This protein carries a polyhistidine tag at the C-terminus.

Conjugate

PE

Excitation Wavelength: 488 nm / 561 nm

Emission Wavelength: 575 nm

Formulation

Lyophilized from 0.22 μm filtered solution in PBS, 0.2% BSA, pH7.4 with trehalose as protectant.

Contact us for customized product form or formulation.

Reconstitution

Please see Certificate of Analysis for specific instructions.

For best performance, we strongly recommend you to follow the reconstitution protocol provided in the CoA.

Storage

For long term storage, the product should be stored at lyophilized state at -20°C or lower.

Please protect from light and avoid repeated freeze-thaw cycles.

This product is stable after storage at:

- 20°C to -70°C for 12 months in lyophilized state;
- 70°C for 3 months under sterile conditions after reconstitution.

Star Staining fluorescent-labeled products are developed by a new-generation site-specific labeling technology with Star Standard quality at ACROBiosystems

★ Using new-generation site-specific labeling technology to maintain natural bioactivity.


★ No non-specific binding to non-transduced PBMCs.

★ High specificity and sensitivity verified by flow cytometry.

★ High homogeneity and high batch-to-batch consistency.

Evaluation of CAR expression

FACS Analysis of TIGIT binding to 293T overexpressing CD155 cells



Flow cytometric analysis of 293T overexpressing CD155 cells staining with PE-Labeled Human TIGIT Protein, His Tag (Cat. No. TIT-HP2H3) at 1:50 dilution (2 μL of the stock solution corresponds to labeling of 5e5 cells in a final volume of 100 μL), compared with negative control protein(Fig. B and A). PE signal was used to



evaluate the binding activity (QC tested).

Background

T-cell immunoreceptor with Ig and ITIM domains (TIGIT) is also known as V-set and immunoglobulin domain-containing protein 9 (VSIG9), V-set and transmembrane domain-containing protein 3 (VSTM3),which belongs to single-pass type I membrane protein containing an immunoglobulin variable domain, a transmembrane domain and an immunoreceptor tyrosine-based inhibitory motif (ITIM). TIGIT is expressed at low levels on peripheral memory and regulatory CD4+ T-cells and NK cells and is up-regulated following activation of these cells (at protein level). TIGIT binds with high affinity to the poliovirus receptor (PVR) which causes increased secretion of IL10 and decreased secretion of IL12B and suppresses T-cell activation by promoting the generation of mature immunoregulatory dendritic cells.

Discounts, Gifts,
and more!

