

Synonym

CD40,Bp50,CDW40,MGC9013,TNFRSF5,p50

Source

Biotinylated Human CD40 Protein, Fc,Avitag(TN5-H82F9) is expressed from human 293 cells (HEK293). It contains AA Glu 21 - Arg 193 (Accession # [P25942-1](#)).

Predicted N-terminus: Glu 21

Molecular Characterization

|                                     |                                 |     |
|-------------------------------------|---------------------------------|-----|
| CD40(Glu 21 - Arg 193 )<br>P25942-1 | Fc(Pro 100 - Lys 330)<br>P01857 | Avi |
|-------------------------------------|---------------------------------|-----|

This protein carries a human IgG1 Fc tag at the C-terminus, followed by an Avi tag (Avitag™).

The protein has a calculated MW of 48.1 kDa. The protein migrates as 60-66 kDa under reducing (R) condition (SDS-PAGE) due to glycosylation.

Labeling

*Biotinylation of this product is performed using Avitag™ technology. Briefly, the single lysine residue in the Avitag is enzymatically labeled with biotin.*

Protein Ratio

Passed as determined by the HABA assay / binding ELISA.

Purity

>95% as determined by SDS-PAGE.

>90% as determined by SEC-MALS.

Formulation

Lyophilized from 0.22 µm filtered solution in Tris with Glycine, Arginine and NaCl, pH7.5 with trehalose as protectant.

Contact us for customized product form or formulation.

Reconstitution

Please see Certificate of Analysis for specific instructions.

*For best performance, we strongly recommend you to follow the reconstitution protocol provided in the CoA.*

Storage


For long term storage, the product should be stored at lyophilized state at -20°C or lower.

*Please avoid repeated freeze-thaw cycles.*

This product is stable after storage at:


- 20°C to -70°C for 12 months in lyophilized state;
- 70°C for 3 months under sterile conditions after reconstitution.

SDS-PAGE



Biotinylated Human CD40 Protein, Fc,Avitag on SDS-PAGE under reducing (R) condition. The gel was stained with Coomassie Blue. The purity of the protein is greater than 95%.

SEC-MALS





The purity of Biotinylated Human CD40 Protein, Fc,Avitag (Cat. No. TN5-H82F9) is more than 90% and the molecular weight of this protein is around 105-120 kDa verified by SEC-MALS.

[Report](#)

Bioactivity-ELISA






Immobilized Human CD40 Ligand, His,Flag Tag, premium grade (Cat. No. CDL-H52Db) at 5 µg/mL (100 µL/well) can bind Biotinylated Human CD40, Fc,Avitag (Cat. No. TN5-H82F9) with a linear range of 0.01-0.625 µg/mL (QC tested).

Immobilized Biotinylated Human CD40, Fc,Avitag (Cat. No. TN5-H82F9) at 5 µg/mL (100 µL/well) on streptavidin precoated (0.5 µg/well) plate, can bind Human CD40 Ligand, His,Flag Tag, premium grade (Cat. No. CDL-H52Db) with a linear range of 0.039-1.25 µg/mL (Routinely tested).

Bioactivity-FACS



FACS analysis shows that Biotinylated Human CD40, Fc,Avitag (Cat. No. TN5-H82F9) can bind to 293T cells overexpressing human CD40L. The concentration of Biotinylated Human CD40 is 0.1 µg/mL (Routinely tested).

Background

CD40 is also known as TNFRSF5, Bp50, CDW40, MGC9013, TNFRSF5 and p50, is a member of the TNF receptor superfamily which are single transmembrane-spanning glycoproteins, and plays an essential role in mediating a broad variety of immune and inflammatory responses including T cell-dependent immunoglobulin class switching, memory B cell development, and germinal center formation. CD40 is a costimulatory protein found on antigen presenting cells and is required for their activation. The binding of CD154 (CD40L) on TH cells to CD40 activates antigen presenting cells and induces a variety of downstream effects. CD40 contains 4 cysteine-rich repeats in the extracellular domain, and is expressed in B cells, dendritic cells, macrophages, endothelial cells, and several tumor cell lines. The extracellular domain has the cysteine-rich repeat regions, which are characteristic for many of the receptors of the TNF superfamily. Interaction of CD40 with its ligand, CD40L, leads to aggregation of CD40 molecules, which in turn interact with cytoplasmic components to initiate signaling pathways. Early studies on the CD40-CD40L system revealed its role in humoral immunity. Defects in CD40 result in hyper-IgM immunodeficiency type 3 (HIGM3), an autosomal recessive disorder characterized by an inability of B cells to undergo isotype switching, as well as an inability to mount an antibody-specific immune response, and a lack of germinal center formation.

