



Abbexa Ltd, Innovation Centre, Cambridge Science Park, Cambridge, CB4 0EY, UK
Telephone: +44 (0) 1223 755950 - Fax: +44 (0) 1223 755951 - E-Mail: info@abbexa.com

Hydroxyacylglutathione Hydrolase (HAGH) Antibody


Catalogue No.: abx005073




Western blot analysis of extracts of various cell lines, using HAGH antibody (abx005073) at 1/1000 dilution.



Immunohistochemistry of paraffin-embedded human liver cancer using HAGH antibody (abx005073) at dilution of 1/100 (40x lens).



Immunohistochemistry of paraffin-embedded human esophageal cancer using HAGH antibody (abx005073) at dilution of 1/100 (40x lens).



Immunofluorescence analysis of A549 cells using HAGH antibody (abx005073). Blue: DAPI for nuclear staining.

HAGH Antibody is a Rabbit Polyclonal antibody against HAGH. The enzyme encoded by this gene is classified as a thiolesterase and is responsible for the hydrolysis of S-lactoyl-glutathione to reduced glutathione and D-lactate. Three transcript variants encoding different isoforms have been found for this gene.

Target: HAGH

Reactivity: Human, Mouse, Rat

Abbexa Ltd, Innovation Centre, Cambridge Science Park, Cambridge, CB4 0EY, UK
Telephone: +44 (0) 1223 755950 - Fax: +44 (0) 1223 755951 - E-Mail: info@abbexa.com

Host:	Rabbit
Clonality:	Polyclonal
Tested Applications:	WB, IHC, IF/ICC
Recommended dilutions:	WB: 1/500 - 1/2000, IHC: 1/50 - 1/200, IF/ICC: 1/10 - 1/100. Optimal dilutions/concentrations should be determined by the end user.
Immunogen:	Recombinant protein of human HAGH.
Purification:	Affinity purified.
Form:	Liquid
Isotype:	IgG
Conjugation:	Unconjugated
Storage:	Aliquot and store at -20 °C. Avoid repeated freeze/thaw cycles.
Molecular Weight:	Calculated MW: 25 kDa/28 kDa/33 kDa Observed MW: 33 kDa
Swiss Prot:	Q16775
GeneID:	3029
Gene Symbol:	HAGH
Concentration:	> 1 mg/ml
Buffer:	PBS, pH 7.3, 0.02% sodium azide, 50% glycerol.
Note:	This product is for research use only.