


## **DATASHEET**

Abbexa Ltd, Innovation Centre, Cambridge Science Park, Cambridge, CB4 0EY, UK Telephone: +44 (0) 1223 755950 - Fax: +44 (0) 1223 755951 - E-Mail: info@abbexa.com

## mouse p21Cip1 (pS125) Antibody

Catalogue No.:abx025059



The protein encoded by this gene belongs to the highly conserved cyclin family, whose members are characterized by a dramatic periodicity in protein abundance through the cell cycle. Cyclins function as regulators of CDK kinases. Different cyclins exhibit distinct expression and degradation patterns which contribute to the temporal coordination of each mitotic event. This cyclin forms a complex with and functions as a regulatory subunit of CDK4 or CDK6, whose activity is required for cell cycle G1/S transition. This protein has been shown to interact with and be involved in the phosphorylation of tumor suppressor protein Rb. The CDK4 activity associated with this cyclin was reported to be necessary for cell cycle progression through G2 phase into mitosis after UV radiation. Several transcript variants encoding different isoforms have been found for this gene.

mouse p21Cip1 (pS125)

Reactivity: Mouse

Host: Rabbit

Clonality: Polyclonal

Tested Applications: DB

Recommended dilutions: Optimal dilutions/concentrations should be determined by the end user.

**Immunogen:** Mouse p21Cip1 (phospho-Ser125).

**Purification:** Peptide Affinity Purified Rabbit Polyclonal Antibody.

**Isotype:** IgG

Conjugation: Unconjugated

Specificity: This mouse p21Cip1 Antibody is generated from rabbits immunized with a KLH conjugated

synthetic phosphopeptide corresponding to amino acid residues surrounding S125 of mouse

p21Cip1.

**Storage:** Aliquot and store at -20 °C. Avoid repeated freeze/thaw cycles.



## **DATASHEET**

Abbexa Ltd, Innovation Centre, Cambridge Science Park, Cambridge, CB4 0EY, UK Telephone: +44 (0) 1223 755950 - Fax: +44 (0) 1223 755951 - E-Mail: info@abbexa.com

Swiss Prot: P39689

Buffer: PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed

by two-step phosphospecific peptide affinity purification.

**Note:** This product is for research use only.