

Abbexa Ltd, Innovation Centre, Cambridge Science Park, Cambridge, CB4 0EY, UK
Telephone: +44 (0) 1223 755950 - Fax: +44 (0) 1223 755951 - E-Mail: info@abbexa.com

Mouse Platelet Endothelial Cell Adhesion Molecule / CD31 (PECAM1) ELISA Kit

Catalogue No.: abx050031



Mouse CD31 ELISA kit is an ELISA kit against Mouse CD31.

Target: Platelet Endothelial Cell Adhesion Molecule / CD31 (PECAM1)

Reactivity: Mouse

Tested Applications: ELISA

Recommended dilutions: Optimal dilutions/concentrations should be determined by the end user.

Test Range: 156 pg/ml - 10000 pg/ml

Sensitivity: < 10 pg/ml

Validity: The validity for this kit is 6 months.

Storage: Store at 2°C to 8°C upon receipt.

Stability: The stability of the kit is determined by the rate of activity loss. The loss rate is less than 5% within the expiration date under appropriate storage conditions. To minimize performance fluctuations, operation procedures and lab conditions should be strictly controlled. It is also strongly suggested that the whole assay is performed by the same user throughout.

Swiss Prot: [Q08481](#)

GeneID: [18613](#)

Gene Symbol: PECAM1

Ensembl: ENSMUSG00000020717

Standard Concentration: 10 ng

Abbexa Ltd, Innovation Centre, Cambridge Science Park, Cambridge, CB4 0EY, UK
Telephone: +44 (0) 1223 755950 - Fax: +44 (0) 1223 755951 - E-Mail: info@abbexa.com

Standard Form: Lyophilized

ELISA Type: Sandwich

ELISA Data: Quantitative

Sample Type: Serum, plasma, tissue lysates, culture media and other biological fluids.

Note: This product is for research use only.
The range and sensitivity is subject to change. Please contact us for the latest product information.
For accurate results, sample concentrations must be diluted to mid-range of the kit. If you require a specific range, please contact us in advance or write your request in your order comments.
Please note that our ELISA and CLIA kits are optimised for detection of native samples, rather than recombinant proteins. We are unable to guarantee detection of recombinant proteins, as they may have different sequences or tertiary structures to the native protein.