



Tachykinin Receptor 2 (TACR2) Antibody


Catalogue No.: abx102297




Western blot analysis of Human Lung lysate, using Human TACR2 Antibody (1 µg/ml) and HRP-conjugated Goat Anti-Rabbit antibody ([abx400043](#), 0.2 µg/ml).




Western blot analysis of Human Hela cell lysate, using Rabbit Anti-Human TACR2 Antibody (1 µg/ml) and HRP-conjugated Goat Anti-Rabbit antibody ([abx400043](#), 0.2 µg/ml).




Western blot analysis of Pig Small intestine lysate, using Rabbit Anti-Human TACR2 Antibody (1 µg/ml) and HRP-conjugated Goat Anti-Rabbit antibody ([abx400043](#), 0.2 µg/ml).




Western blot analysis of Porcine Heart lysate, using Rabbit Anti-Human TACR2 Antibody (1 µg/ml) and HRP-conjugated Goat Anti-Rabbit antibody ([abx400043](#), 0.2 µg/ml).



Western blot analysis of the recombinant protein.



IHC-P analysis of Hela Cells, with FITC staining.



IHC-P analysis of Human Pancreatic Cancer Tissue, with DAB staining.

Polyclonal Antibody to Tachykinin Receptor 2 (TACR2).

Target: Tachykinin Receptor 2 (TACR2)

Reactivity: Human

Host: Rabbit

Clonality: Polyclonal

Tested Applications: WB, IHC, IF/ICC, IP

Recommended dilutions: Optimal dilutions/concentrations should be determined by the end user.

Immunogen: TACR2 (Val92-Leu290)

Purification: Antigen-specific affinity chromatography followed by Protein A affinity chromatography.

Form: Liquid

Conjugation: Unconjugated

Storage: Store at 4 °C for frequent use. Aliquot and store at -20 °C for one year. Avoid repeated freeze/thaw cycles.

Buffer: PBS, pH 7.4, containing 0.01% Sarcosyl, 1 mM DTT, 5% Trehalose and Proclin300.

Note: This product is for research use only.