



Abbexa Ltd, Innovation Centre, Cambridge Science Park, Cambridge, CB4 0EY, UK
Telephone: +44 (0) 1223 755950 - Fax: +44 (0) 1223 755951 - E-Mail: info@abbexa.com

Histone Cluster 1, H2ag (HIST1H2AG) Antibody


Catalogue No.: abx102875



Western blot analysis of recombinant Human HIST1H2AG.



IHC-P analysis of Human Stomach Tissue, with DAB staining.



IHC-P analysis of Human Prostate Gland Tissue, with DAB staining.

Polyclonal Antibody to Histone Cluster 1, H2ag (HIST1H2AG).

Target: Histone Cluster 1, H2ag (HIST1H2AG)

Reactivity: Human

Host: Rabbit

Clonality: Polyclonal

Tested Applications: WB, IHC, IF/ICC, IP

Recommended dilutions: WB: 0.5 - 2 µg/ml, IHC: 5 - 20 µg/ml, ICC: 5 - 20 µg/ml. Optimal dilutions/concentrations should be determined by the end user.

Immunogen: HIST1H2AG (Met1-Lys130)

Abbexa Ltd, Innovation Centre, Cambridge Science Park, Cambridge, CB4 0EY, UK
Telephone: +44 (0) 1223 755950 - Fax: +44 (0) 1223 755951 - E-Mail: info@abbexa.com

Purification:	Antigen-specific affinity chromatography followed by Protein A affinity chromatography.
Form:	Liquid
Conjugation:	Unconjugated
Storage:	Store at 4 °C for frequent use. Aliquot and store at -20 °C for one year. Avoid repeated freeze/thaw cycles.
Buffer:	0.01 M PBS, pH 7.4, containing 0.05% Proclin-300, 50% glycerol.
Note:	This product is for research use only.