



Abbexa Ltd, Innovation Centre, Cambridge Science Park, Cambridge, CB4 0EY, UK
Telephone: +44 (0) 1223 755950 - Fax: +44 (0) 1223 755951 - E-Mail: info@abbexa.com

Endoplasmic Reticulum To Nucleus Signalling 1 (ERN1) Antibody


Catalogue No.: abx128006



Western blot analysis of recombinant Human ERN1.



Western Blot analysis of (1) Human K562 cell lysate (2) Human 293T cell lysate, and (3) Monkey COS7 cell lysate, using Rabbit Anti-Human ERN1 Antibody (2 µg/ml) and HRP-conjugated Goat Anti-Rabbit antibody ([abx400043](#), 0.2 µg/ml).



IHC-P analysis of Pig Cerebrum Tissue, with DAB staining, using Rabbit Anti-Human ERN1 Antibody (10 µg/ml) and HRP-conjugated Goat Anti-Rabbit antibody ([abx400043](#), 2 µg/ml).

Endoplasmic Reticulum To Nucleus Signalling 1 Antibody is a Rabbit Polyclonal against Endoplasmic Reticulum To Nucleus Signalling 1.

Target: Endoplasmic Reticulum To Nucleus Signalling 1 (ERN1)

Reactivity: Human

Host: Rabbit

Clonality: Polyclonal

Tested Applications: WB, IHC, IF/ICC, IP

Recommended dilutions: WB: 0.5-2 µg/ml, IHC: 0.5-2 µg/ml, ICC: 0.5-2 µg/ml. Optimal dilutions/concentrations should be determined by the end user.

Abbexa Ltd, Innovation Centre, Cambridge Science Park, Cambridge, CB4 0EY, UK
Telephone: +44 (0) 1223 755950 - Fax: +44 (0) 1223 755951 - E-Mail: info@abbexa.com

Immunogen:	Recombinant ERN1 (Phe571-Phe832) expressed in E. coli.
Purification:	Antigen-specific affinity chromatography followed by Protein A affinity chromatography.
Form:	Liquid
Conjugation:	Unconjugated
Storage:	Store at 4 °C for frequent use. Aliquot and store at -20 °C for one year. Avoid repeated freeze/thaw cycles.
Swiss Prot:	<u>O75460</u>
Buffer:	0.01 M PBS, pH 7.4, containing 0.05% Proclin-300, 50% glycerol.
Note:	This product is for research use only.