



Abbexa Ltd, Innovation Centre, Cambridge Science Park, Cambridge, CB4 0EY, UK
Telephone: +44 (0) 1223 755950 - Fax: +44 (0) 1223 755951 - E-Mail: info@abbexa.com

High Mobility Group Protein 17 (HMG17) Antibody


Catalogue No.: abx128751




Western blot analysis of (1) Human Jurkat Cells, (2) Human K562 Cells, (3) Human HL-60 Cells and (4) Human HeLa cells.




Western blot analysis of recombinant Human HMG17.



IHC-P analysis of Human Stomach Tissue, with DAB staining.




IHC-P analysis of Human Kidney Tissue, with DAB staining.



IHC-P analysis of Human Stomach Cancer Tissue, with DAB staining.

Abbexa Ltd, Innovation Centre, Cambridge Science Park, Cambridge, CB4 0EY, UK
Telephone: +44 (0) 1223 755950 - Fax: +44 (0) 1223 755951 - E-Mail: info@abbexa.com



IHC-P analysis of Human Skin Cancer Tissue, with DAB staining.

High Mobility Group Protein 17 Antibody is a Rabbit Polyclonal against High Mobility Group Protein 17.

Target: High Mobility Group Protein 17 (HMG17)

Reactivity: Human

Host: Rabbit

Clonality: Polyclonal

Tested Applications: WB, IHC, IF/ICC, IP

Recommended dilutions: Optimal dilutions/concentrations should be determined by the end user.

Immunogen: HMG17 (Pro2-Gln81)

Purification: Antigen-specific affinity chromatography followed by Protein A affinity chromatography.

Form: Liquid

Conjugation: Unconjugated

Storage: Store at 4 °C for frequent use. Aliquot and store at -20 °C for one year. Avoid repeated freeze/thaw cycles.

Buffer: 0.01 M PBS, pH 7.4, containing 0.05% Proclin-300, 50% glycerol.

Note: This product is for research use only.