


Abbexa Ltd, Innovation Centre, Cambridge Science Park, Cambridge, CB4 0EY, UK Telephone: +44 (0) 1223 755950 - Fax: +44 (0) 1223 755951 - E-Mail: info@abbexa.com


## DATASHEET

## Branched Chain Alpha-Ketoacid Dehydrogenase Kinase (BCKDK) Antibody

Catalogue No.:abx130391




Western blot analysis of (1) Rat Serum, (2) Rat Pancreas Tissue, (3) Rat Brain Tissue and (4) Rat Adrenal Gland Tissue.




Western blot analysis of recombinant Rat BCKDK.



IHC-P analysis of Rat Skeletal Muscle Tissue, with DAB staining.



IHC-P analysis of Rat Stomach Tissue, with DAB staining.




IHC-P analysis of Rat Heart Tissue, with DAB staining.



## Abbexa Ltd, Innovation Centre, Cambridge Science Park, Cambridge, CB4 0EY, UK Telephone: +44 (0) 1223 755950 - Fax: +44 (0) 1223 755951 - E-Mail: info@abbexa.com

## DATASHEET



IHC-P analysis of Rat Liver Tissue, with DAB staining.

Branched Chain Alpha-Ketoacid Dehydrogenase Kinase Antibody is a Rabbit Polyclonal against Branched Chain Alpha-Ketoacid Dehydrogenase Kinase.

Target:	Branched Chain Alpha-Ketoacid Dehydrogenase Kinase (BCKDK)
Reactivity:	Rat
Host:	Rabbit
Clonality:	Polyclonal
Tested Applications:	WB, IHC, IF/ICC, IP
Recommended dilutions	: Optimal dilutions/concentrations should be determined by the end user.
Immunogen:	BCKDK (Arg44-Thr223)
Purification:	Antigen-specific affinity chromatography followed by Protein A affinity chromatography.
Form:	Liquid
Conjugation:	Unconjugated
Storage:	Store at 4 °C for frequent use. Aliquot and store at -20 °C for one year. Avoid repeated freeze/thaw cycles.
Swiss Prot:	<u>Q00972</u>
Buffer:	0.01 M PBS, pH 7.4, containing 0.05% Proclin-300, 50% glycerol.
Note:	This product is for research use only.