





Abbexa Ltd, Innovation Centre, Cambridge Science Park, Cambridge, CB4 0EY, UK Telephone: +44 (0) 1223 755950 - Fax: +44 (0) 1223 755951 - E-Mail: info@abbexa.com

Human Procollagen I N-Terminal Propeptide (PINP) Peptide

Catalogue No.:abx652144



SDS-PAGE analysis of recombinant Human PINP.



Gene sequencing extract of Human PINP,

Procollagen I N-Terminal Propeptide (PINP) Protein is a Recombinant Human protein expressed in E. coli.

This protein is the immunogen for the following antibodies: <u>abx103755</u>

Target: Procollagen I N-Terminal Propeptide (PINP)

Origin: Human

Host: E. coli

Tested Applications: WB, SDS-PAGE

Purity: > 90%

Form: Lyophilized

Reconstitution: Reconstitute in 100 mM NaHCO₃, 500 mM NaCl, pH 8.3, to a concentration of 0.1-1.0 mg/ml. Do not

vortex.

Conjugation: Unconjugated

Storage: Store at 2-8 °C for up to one month. Store at -80 °C for up to one year. Avoid repeated freeze/thaw

cycles.

Expression: Recombinant



DATASHEET

Abbexa Ltd, Innovation Centre, Cambridge Science Park, Cambridge, CB4 0EY, UK Telephone: +44 (0) 1223 755950 - Fax: +44 (0) 1223 755951 - E-Mail: info@abbexa.com

Molecular Weight: Calculated MW: 27.9 kDa

Observed MW (SDS-PAGE): 40 kDa

Possible reasons why the actual band size differs from the predicted band size:

- 1. Splice variants. Alternative splicing may create different sized proteins from the same gene.
- 2. Relative charge. The composition of amino acids may affect the charge of the protein.
- 3. Post-translational modification. Phosphorylation, glycoslyation, methylation etc. may affect the band
- 4. Post-translational cleavage. Many proteins are synthesised as pro-proteins, and then cleaved to give the active form.
- 5. Polymerisation of the target protein. Dimerisation, multimerisation etc. will increase the band size observed.

Swiss Prot: P02452

Tag: N-terminal His tag

Activity: Not tested

Concentration: Prior to lyophilization: 200 μg/ml

Buffer: Prior to lyophilization: 100 mM NaHCO₃, 500 mM NaCl, pH 8.3, containing 0.01% Sarcosyl, 5%

Trehalose.

Note: This product is for research use only.