

**Anti-Human IL-17D Antibody**  
**Catalog # ABG10225****Specification**

---

**Anti-Human IL-17D Antibody - Product Information**

Application	<b>WB, IHC, E</b>
Reactivity	<b>Human</b>
Host	<b>Rabbit</b>
Clonality	<b>Polyclonal</b>

**Anti-Human IL-17D Antibody - Additional Information****Preparation**

Produced from sera of rabbits immunized with highly pure recombinant Human IL-17D. Anti-Human IL-17D specific antibody was purified by affinity chromatography employing an immobilized Human IL-17D matrix.

**WesternBlot**

To detect Human IL-17D by Western Blot analysis this antibody can be used at a concentration of 0.1 - 0.2 µg/ml. Used in conjunction with compatible secondary reagents the detection limit for recombinant Human IL-17D is 1.5 - 3.0 ng/lane, under either reducing or non-reducing conditions.

**Sandwich**

To detect Human IL-17D by sandwich ELISA (using 100µl/well antibody solution) a concentration of 0.5 - 2.0 µg/ml of this antibody is required. This antigen affinity purified antibody, in conjunction with BioGems's Biotinylated Anti-Human IL-17D (60-017DBT) as a detection antibody, allows the detection of at least 0.2 - 0.4 ng/well of recombinant Human IL-17D.

**Immunohistochemistry**

This antibody stained formalin-fixed, paraffin-embedded sections of human normal pancreas. The recommended concentration is 2.5 µg/mL for two hours at room temperature. An HRP-labeled polymer detection system was used with a

DAB chromogen. Optimal results for these conditions were achieved without antigen retrieval. Optimal concentrations and conditions may vary. <br /> <br /> <em><span style="font-size: 10px;">Information and photo are courtesy of Flagship Biosciences, LLC - Histology Services. </span></em>

**Formulation**

A sterile filtered antibody solution was lyophilized from PBS, pH 7.2.

**Reconstitution**

Centrifuge vial prior to opening.  
Reconstitute in sterile water to a concentration of 0.1-1.0 mg/ml.

**Storage**

-20°C

**Precautions**

Anti-Human IL-17D Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

**Anti-Human IL-17D Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)