

Anti-Human IL-8 Antibody
Catalog # ABG10308**Specification**

Anti-Human IL-8 Antibody - Product Information

Application	WB, IHC, E
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal

Anti-Human IL-8 Antibody - Additional Information**Preparation**

Produced from sera of rabbits pre-immunized with highly pure recombinant Human IL-8. Anti-Human IL-8 specific antibody was purified by affinity chromatography employing immobilized Human IL-8 matrix.

WesternBlot

To detect Human IL-8 by Western Blot analysis this antibody can be used at a concentration of 0.1 - 0.2 µg/ml. When used in conjunction with compatible secondary reagents, the detection limit for recombinant Human IL-8 is 1.5 - 3.0 ng/lane, under either reducing or non-reducing conditions.

Sandwich

To detect Human IL-8 by sandwich ELISA (using 100 µl/well antibody solution) a concentration of 0.5 - 2.0 µg/ml of this antibody is required. This antigen affinity purified antibody, in conjunction with BioGems's Biotinylated Anti-Human IL-8 (60-008BT) as a detection antibody, allows the detection of at least 0.2 - 0.4 ng/well of recombinant Human IL-8.

Immunohistochemistry

This antibody stained formalin-fixed, paraffin-embedded sections of normal human skin. The recommended concentrations are 1.0 µg/ml for two hours at room temperature or 0.25 µg/ml overnight at 4°C. An HRP-labeled polymer detection system was used with a non-alcohol soluble AEC chromogen and a

proteinase K antigen retrieval. Optimal concentrations and conditions may vary.
 *Additional Immunostaining data available. Please contact Tech Support for information.

Neutralization

To yield one-half maximal inhibition [**ND**₅₀] of the biological activity of Human IL-8 (100ng/ml), a concentration of 2.0 - 3.0 µg/ml of this antibody is required.

Formulation

A sterile filtered antibody solution was lyophilized from PBS, pH 7.2.

Reconstitution

Centrifuge vial prior to opening. Reconstitute in sterile water to a concentration of 0.1-1.0 mg/ml.

Storage

-20°C

Precautions

Anti-Human IL-8 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Anti-Human IL-8 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)