

**Anti-BMPR2 Picoband Antibody**  
**Catalog # ABO10056****Specification****Anti-BMPR2 Picoband Antibody - Product Information**

Application	<b>WB, E</b>
Primary Accession	<a href="#">Q13873</a>
Host	<b>Rabbit</b>
Reactivity	<b>Human, Mouse, Rat</b>
Clonality	<b>Polyclonal</b>
Format	<b>Lyophilized</b>

**Description**

Rabbit IgG polyclonal antibody for BMPR2 detection. Tested with WB, Direct ELISA in Human;Mouse;Rat.

**Reconstitution**

Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

**Anti-BMPR2 Picoband Antibody - Additional Information**

**Gene ID 659**

**Other Names**

Bone morphogenetic protein receptor type-2, BMP type-2 receptor, BMPR-2, 2.7.11.30, Bone morphogenetic protein receptor type II, BMP type II receptor, BMPR-II, BMPR2, PPH1

**Application Details**

Western blot, 0.1-0.5 µg/ml  
Direct ELISA, 0.1-0.5 µg/ml

**Subcellular Localization**

Cell membrane.

**Tissue Specificity**

Highly expressed in heart and liver.

**Contents**

Each vial contains 4mg Trehalose, 0.9mg NaCl, 0.2mg Na<sub>2</sub>HPO<sub>4</sub>, 0.05mg NaN<sub>3</sub>.

**Anti-BMPR2 Picoband Antibody - Background**

Bone morphogenetic protein receptor type II or BMPR2 is a serine/threonine receptor kinase. This gene encodes a member of the bone morphogenetic protein (BMP) receptor family of transmembrane serine/threonine kinases. The ligands of this receptor are BMPs, which are members of the TGF-beta superfamily. BMPs are involved in endochondral bone formation and embryogenesis. These proteins transduce their signals through the formation of heteromeric complexes of two different types of serine (threonine) kinase receptors: type I receptors of about 50-55 kD and type II receptors of about 70-80 kD. Type II receptors bind ligands in the absence of type I receptors, but they require their respective type I receptors for signaling, whereas type I receptors require their respective type II receptors for ligand binding. Mutations in this gene have been associated with primary pulmonary hypertension, both familial and fenfluramine-associated, and with pulmonary venoocclusive disease.

**Immunogen**

E. coli-derived human BMPR2 recombinant protein (Position: R455-K512).

**Cross Reactivity**

No cross reactivity with other proteins.

**Storage**

**At -20°C; for one year. After reconstitution, at 4°C; for one month. It can also be aliquotted and stored frozen at -20°C; for a longer time. Avoid repeated freezing and thawing.**

**Anti-BMPR2 Picoband Antibody - Protein Information**

**Name** BMPR2

**Synonyms** PPH1

**Function**

On ligand binding, forms a receptor complex consisting of two type II and two type I transmembrane serine/threonine kinases. Type II receptors phosphorylate and activate type I receptors which autophosphorylate, then bind and activate SMAD transcriptional regulators. Binds to BMP7, BMP2 and, less efficiently, BMP4. Binding is weak but enhanced by the presence of type I receptors for BMPs. Mediates induction of adipogenesis by GDF6.

**Cellular Location**

Cell membrane; Single-pass type I membrane protein

**Tissue Location**

Highly expressed in heart and liver.

**Anti-BMPR2 Picoband Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)

- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)