

## Anti-CHX10 Picoband Antibody

Catalog # ABO13045

### Specification

#### Anti-CHX10 Picoband Antibody - Product Information

Application	WB
Primary Accession	<a href="#">P58304</a>
Host	Rabbit
Reactivity	Human, Mouse
Clonality	Polyclonal
Format	Lyophilized

#### Description

Rabbit IgG polyclonal antibody for Visual system homeobox 2(VSX2) detection. Tested with WB in Human;Mouse.

#### Reconstitution

Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

#### Anti-CHX10 Picoband Antibody - Additional Information

Gene ID 338917

#### Other Names

Visual system homeobox 2, Ceh-10 homeodomain-containing homolog, Homeobox protein CHX10, VSX2, CHX10, HOX10

#### Calculated MW

39411 MW KDa

#### Application Details

Western blot, 0.1-0.5 µg/ml, Mouse, Human<br>

#### Subcellular Localization

Nucleus.

#### Tissue Specificity

Abundantly expressed in retinal neuroblasts during eye development and in the inner nuclear layer of the adult retina. Within this layer, expression is stronger in the outer margin where bipolar cells predominate.

#### Contents




Figure 1. Western blot analysis of CHX10 using anti-CHX10 antibody (ABO13045).

#### Anti-CHX10 Picoband Antibody - Background

Visual system homeobox 2 is a protein that in humans is encoded by the VSX2 gene. This gene encodes a homeobox protein originally described as a retina-specific transcription factor. Mutations in this gene are associated with microphthalmia, cataracts and iris abnormalities. In situ hybridization to human fetal retinal sections detected CHX10 expression in retinal neuroblasts at all stages examined. Human CHX10 is expressed in progenitor cells of the developing neuroretina and in the inner nuclear layer of the mature retina.

Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na<sub>2</sub>HPO<sub>4</sub>, 0.05mg NaN<sub>3</sub>.

**Immunogen**

E. coli-derived human CHX10 recombinant protein (Position: A249-A361). Human CHX10 shares 90.3% amino acid (aa) sequence identity with mouse CHX10.

**Purification**

Immunogen affinity purified.

**Cross Reactivity**

No cross reactivity with other proteins.

**Storage**

**At -20°C for one year. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for a longer time. Avoid repeated freezing and thawing.**

**Anti-CHX10 Picoband Antibody - Protein Information****Name VSX2****Synonyms CHX10, HOX10****Function**

Acts as a transcriptional regulator through binding to DNA at the consensus sequence 5'-(TC)TAATT[AG][AG]-3' upstream of gene promoters (PubMed:<a href="http://www.uniprot.org/citations/27301076" target="\_blank">27301076</a>). Plays a significant role in the specification and morphogenesis of the sensory retina (By similarity). Mediates differentiation of V2a interneurons by repression of motor neuron gene transcription, via competitively binding to response elements that are activated by the ISL1-LHX3 complex, such as VSX1 (PubMed:<a href="http://www.uniprot.org/citations/17919464" target="\_blank">17919464</a>, PubMed:<a href="http://www.uniprot.org/citations/27477290" target="\_blank">27477290</a>). Acts as a positive transcriptional regulator of NXNL1; regulation is significantly increased in

synergy with VSX1 (By similarity). Acts as a negative transcriptional regulator of MITF (By similarity). Represses SAG transcription by competitive inhibition of ISL1-LHX3 response elements (PubMed:<a href="http://www.uniprot.org/citations/16236706" target="\_blank">16236706</a>, PubMed:<a href="http://www.uniprot.org/citations/27477290" target="\_blank">27477290</a>). Binds to the photoreceptor conserved element-1 (PCE- 1) in the promoter of rod photoreceptor arrestin SAG and acts as a transcriptional repressor (By similarity). Plays a significant role in the specification and morphogenesis of the sensory retina (By similarity). Involved in the development of retinal ganglion cells (RGCs) which leads to release of SHH by RGCs, promoting Hedgehog signaling and subsequent proliferation of retinal progenitor cells (By similarity). Participates in the development of the cells of the inner nuclear layer, by promoting postnatal differentiation of bipolar cells with a comparable inhibition of rod cell differentiation (By similarity). May play a role in the maintenance of neural retina identity during development by regulation of canonical Wnt genes and CTNNB1 localization, suggesting a role in the regulation of canonical Wnt signaling (PubMed:<a href="http://www.uniprot.org/citations/27301076" target="\_blank">27301076</a>).

#### **Cellular Location**

Nucleus

{ECO:0000250|UniProtKB:Q61412}.

#### **Tissue Location**

Abundantly expressed in retinal neuroblasts during eye development and in the inner nuclear layer of the adult retina Within this layer, expression is stronger in the outer margin where bipolar cells predominate

### **Anti-CHX10 Picoband Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)

- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)