



## Peroxiredoxin VI Antibody (4A3)

Mouse Monoclonal Antibody Catalog # ABV11156

## **Specification**

Peroxiredoxin VI Antibody (4A3) - Product Information

Application WB, E Primary Accession P30041

Reactivity Human, Mouse,

Rat

Host Mouse
Clonality Monoclonal
Isotype Mouse IgG 1

Calculated MW 25035

Peroxiredoxin VI Antibody (4A3) - Additional Information

**Gene ID 9588** 

Positive Control WB analysis:

HeLa, Mouse lung, Rat lung lysates. IHC staining: paraffin embedded human

kidney

Application & Usage Western blot:

1:2000, ELISA.

**Other Names** 

Peroxiredoxin 6, 1-Cys, aiPLA2, AOP2,

NSGPx, p29, PRX.

Target/Specificity

Peroxiredoxin VI

**Antibody Form** 

Liquid

**Appearance** 

Colorless liquid

**Formulation** 

 $100 \mu l$  of antibody in HEPES with 0.15 M NaCl, 0.01 % BSA, 0.03 % sodium azide,

and 50 % glycerol

Handling

The antibody solution should be gently

mixed before use.

# Peroxiredoxin VI Antibody (4A3) - Background

Peroxiredoxin (Prx) is a growing peroxidase family, whose mammalian members have been known to connect with cell proliferation, differentiation, and apoptosis. Many isoforms (about 50 proteins), collected in accordance to the amino acid sequence homology, particularly amino-terminal region containing active site cysteine residue, and the thiol-specific antioxidant activity, distribute throughout all the kingdoms. Among them, mammalian Prx consists of 6 different members grouped into typical 2-Cys, atypical 2-Cys Prx, and 1-Cys Prx. Except Prx VI belonging to 1-Cys Prx subgroup, the other five 2-Cys Prx isotypes have the thioredoxin-dependent peroxidase (TPx) activity utilizing thioredoxin, thioredoxin reductase, and NADPH as a reducing system. Mammalian Prxs are 20 - 30 kDa in molecular size and vary in subcellular localization: Prx I, II, and VI in cytosol, Prx III in mitochondria, Prx IV in ER and secretion, Prx V showing complicated distribution including peroxisome, mitochondria and cytosol. Prx VI is involved in redox regulation of the cell. Can reduce H2O2 and short chain organic, fatty acid, and phospholipid hydroperoxides. May play a role in the regulation of phospholipid turnover as well as in protection against oxidative injury.



## Reconstitution & Storage -20 °C

### **Background Descriptions**

#### **Precautions**

Peroxiredoxin VI Antibody (4A3) is for research use only and not for use in diagnostic or therapeutic procedures.


Peroxiredoxin VI Antibody (4A3) - Protein Information

Name PRDX6

Synonyms AOP2, KIAA0106

#### **Function**

Thiol-specific peroxidase that catalyzes the reduction of hydrogen peroxide and organic hydroperoxides to water and alcohols, respectively (PubMed:<a href="http://www. uniprot.org/citations/9497358" target=" blank">9497358</a>, PubMed:<a href="http://www.uniprot.org/ci tations/10893423" target=" blank">10893423</a>). Can reduce H(2)O(2) and short chain organic, fatty acid, and phospholipid hydroperoxides (PubMed:<a href="http://www.uniprot.org/c itations/10893423" target=" blank">10893423</a>). Also has phospholipase activity, can therefore either reduce the oxidized sn-2 fatty acyl group of phospholipids (peroxidase activity) or hydrolyze the sn-2 ester bond of phospholipids (phospholipase activity) (PubMed:<a href="http://www.uniprot.org/c itations/10893423" target=" blank">10893423</a>, PubMed:<a href="http://www.uniprot.org/ci tations/26830860" target=" blank">26830860</a>). These activities are dependent on binding to phospholipids at acidic pH and to oxidized phospholipds at cytosolic pH (PubMed: <a hr ef="http://www.uniprot.org/citations/10893" 423" target=" blank">10893423</a>). Plays a role in cell protection against oxidative stress by detoxifying peroxides and in phospholipid homeostasis (PubMed:<a href="http://www.uniprot.org/c itations/10893423" target=" blank">10893423</a>). Exhibits acyl-CoA-dependent lysophospholipid





acyltransferase which mediates the conversion of lysophosphatidylcholine (1-acyl-sn-glycero-3- phosphocholine or LPC) into phosphatidylcholine (1,2-diacyl-sn-glycero- 3-phosphocholine or PC) (PubMed:<a href="http://www.uniprot.org/citations/26830860" target="\_blank">26830860" target="\_blank">26830860</a>). Shows a clear preference for LPC as the lysophospholipid and for palmitoyl CoA as the fatty acyl substrate (PubMed:<a href="http://www.uniprot.org/citations/26830860" target="\_blank">26830860</a>).

#### **Cellular Location**

Cytoplasm. Lysosome {ECO:0000250|UniProtKB:O35244}. Note=Also found in lung secretory organelles (lamellar bodies). {ECO:0000250|UniProtKB:O35244}

## Peroxiredoxin VI Antibody (4A3) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture