

Goat Anti-APOA4 Antibody
Peptide-affinity purified goat antibody
Catalog # AF1074a

Specification

Goat Anti-APOA4 Antibody - Product Information

Application	WB, IHC
Primary Accession	P06727
Other Accession	NP_000473 , 337
Reactivity	Human
Host	Goat
Clonality	Polyclonal
Concentration	0.5mg/ml
Isotype	IgG
Calculated MW	45372

Goat Anti-APOA4 Antibody - Additional Information

Gene ID 337

Other Names

Apolipoprotein A-IV, Apo-AIV, ApoA-IV, Apolipoprotein A4, APOA4

Format

0.5 mg IgG/ml in Tris saline (20mM Tris pH7.3, 150mM NaCl), 0.02% sodium azide, with 0.5% bovine serum albumin

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

Goat Anti-APOA4 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Goat Anti-APOA4 Antibody - Protein Information


Name APOA4

Function

May have a role in chylomicrons and VLDL secretion and catabolism. Required for



AF1074a (0.01 µg/ml) staining of Human Duodenum lysate (35 µg protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.



AF1074a (2 µg/ml) staining of paraffin embedded Human Intestine. Steamed antigen retrieval with citrate buffer pH 6, HRP-staining.

Goat Anti-APOA4 Antibody - Background

Apolipoprotein (apo) A-IV gene contains 3 exons separated by two introns. A sequence polymorphism has been identified in the 3'UTR of the third exon. The primary translation product is a 396-residue preprotein which after proteolytic processing is secreted its primary

efficient activation of lipoprotein lipase by ApoC-II; potent activator of LCAT. ApoA-IV is a major component of HDL and chylomicrons.

Cellular Location

Secreted.

Tissue Location

Synthesized primarily in the intestine and secreted in plasma

Goat Anti-APOA4 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

site of synthesis, the intestine, in association with chylomicron particles. Although its precise function is not known, apo A-IV is a potent activator of lecithin-cholesterol acyltransferase in vitro.

Goat Anti-APOA4 Antibody - References

The APOA1/C3/A4/A5 cluster and markers of allostatic load in the Boston Puerto Rican Health Study. Mattei J, et al. Nutr Metab Cardiovasc Dis, 2010 Jul 29. PMID 20674306. Variation at the NFATC2 Locus Increases the Risk of Thiazolidinedione-Induced Edema in the Diabetes REduction Assessment with ramipril and rosiglitazone Medication (DREAM) Study. Bailey SD, et al. Diabetes Care, 2010 Jul 13. PMID 20628086. Physiogenomic analysis of statin-treated patients: domain-specific counter effects within the ACACB gene on low-density lipoprotein cholesterol? Ruaño G, et al. Pharmacogenomics, 2010 Jul. PMID 20602615. No interaction between alcohol consumption and HDL-related genes on HDL cholesterol levels. Marques-Vidal P, et al. Atherosclerosis, 2010 Aug. PMID 20430392. APOA1 and APOA4 gene polymorphisms influence the effects of dietary fat on LDL particle size and oxidation in healthy young adults. Gomez P, et al. J Nutr, 2010 Apr. PMID 20164363.