

Goat Anti-TGM2 Antibody
Peptide-affinity purified goat antibody
Catalog # AF2083a

Specification

Goat Anti-TGM2 Antibody - Product Information

Application	WB
Primary Accession	P21980
Other Accession	NP_004604 , 7052
Reactivity	Human
Host	Goat
Clonality	Polyclonal
Concentration	100ug/200ul
Isotype	IgG
Calculated MW	77329

Goat Anti-TGM2 Antibody - Additional Information

Gene ID 7052

Other Names

Protein-glutamine
gamma-glutamyltransferase 2, 2.3.2.13,
Tissue transglutaminase, Transglutaminase
C, TG(C), TGC, TGase C, Transglutaminase
H, TGase H, Transglutaminase-2, TGase-2,
TGM2

Format

0.5 mg IgG/ml in Tris saline (20mM Tris
pH7.3, 150mM NaCl), 0.02% sodium azide,
with 0.5% bovine serum albumin

Storage

Maintain refrigerated at 2-8°C for up to 6
months. For long term storage store at
-20°C in small aliquots to prevent
freeze-thaw cycles.

Precautions

Goat Anti-TGM2 Antibody is for research use
only and not for use in diagnostic or
therapeutic procedures.

Goat Anti-TGM2 Antibody - Protein Information

Name TGM2

{ECO:0000303|PubMed:17939176,



AF2083a (0.03 µg/ml) staining of Human
Lung lysate (35 µg protein in RIPA buffer).
Primary incubation was 1 hour. Detected by
chemiluminescence.



AF2083a (0.1 µg/ml) staining of A549 lysate
(35 µg protein in RIPA buffer). Primary
incubation was 1 hour. Detected by
chemiluminescence.

Goat Anti-TGM2 Antibody - Background

Transglutaminases are enzymes that catalyze
the crosslinking of proteins by epsilon-gamma
glutamyl lysine isopeptide bonds. While the
primary structure of transglutaminases is not
conserved, they all have the same amino acid
sequence at their active sites and their activity

ECO:0000312|HGNC:HGNC:11778}

Function

Calcium-dependent acyltransferase that catalyzes the formation of covalent bonds between peptide-bound glutamine and various primary amines, such as gamma-amino group of peptide-bound lysine, or mono- and polyamines, thereby producing cross-linked or aminated proteins, respectively (PubMed:9252372, PubMed:23941696, PubMed:31991788). Involved in many biological processes, such as bone development, angiogenesis, wound healing, cellular differentiation, chromatin modification and apoptosis (PubMed:1683874, PubMed:7935379, PubMed:9252372, PubMed:27270573). Acts as a protein-glutamine gamma-glutamyltransferase by mediating the cross-linking of proteins, such as ACO2, HSPB6, FN1, HMGB1, RAP1GDS1, SLC25A4/ANT1, SPP1 and WDR54 (PubMed:23941696, PubMed:24349085, PubMed:29618516, PubMed:30458214). Under physiological conditions, the protein cross-linking activity is inhibited by GTP; inhibition is relieved by Ca(2+) in response to various stresses (PubMed:7649299,

is calcium-dependent. The protein encoded by this gene acts as a monomer, is induced by retinoic acid, and appears to be involved in apoptosis. Finally, the encoded protein is the autoantigen implicated in celiac disease. Two transcript variants encoding different isoforms have been found for this gene.

Goat Anti-TGM2 Antibody - References

I-kappaBalpha depletion by transglutaminase 2 and mu-calpain occurs in parallel with the ubiquitin-proteasome pathway. Kim DS, et al. Biochem Biophys Res Commun, 2010 Aug 20. PMID 20659425.
Redox regulation of transglutaminase 2 activity. Stamnaes J, et al. J Biol Chem, 2010 Aug 13. PMID 20547769.
Decreased S-nitrosylation of tissue transglutaminase contributes to age-related increases in vascular stiffness. Santhanam L, et al. Circ Res, 2010 Jul 9. PMID 20489165.
Coeliac disease autoantibodies mediate significant inhibition of tissue transglutaminase. Byrne G, et al. Clin Immunol, 2010 Sep. PMID 20488756.
Inhibition of tissue transglutaminase sensitizes TRAIL-resistant lung cancer cells through upregulation of death receptor 5. Frese-Schaper M, et al. FEBS Lett, 2010 Jul 2. PMID 20450916.

PubMed:7592956, PubMed:18092889). When secreted, catalyzes cross-linking of proteins of the extracellular matrix, such as FN1 and SPP1 resulting in the formation of scaffolds (PubMed:12506096). Plays a key role during apoptosis, both by (1) promoting the cross-linking of cytoskeletal proteins resulting in condensation of the cytoplasm, and by (2) mediating cross-linking proteins of the extracellular matrix, resulting in the irreversible formation of scaffolds that stabilize the integrity of the dying cells before their clearance by phagocytosis, thereby preventing the leakage of harmful intracellular components (PubMed:7935379, PubMed:9252372). In addition to protein cross-linking, can use different monoamine substrates to catalyze a vast array of protein post-translational modifications: mediates aminylation of serotonin, dopamine, noradrenaline or histamine into glutamine residues of target proteins to generate protein serotonylation, dopaminylation, noradrenalinylation or histaminylation, respectively (PubMed:23797785, PubMed:30867594). Mediates protein serotonylation of small GTPases during activation and aggregation of platelets, leading to constitutive activation of these GTPases (By similarity). Plays a key role in chromatin organization by mediating serotonylation and dopaminylation of histone H3 (PubMed:30867594, PubMed:32273471). Catalyzes serotonylation of 'Gln-5' of histone H3 (H3Q5ser) during serotonergic neuron differentiation, thereby facilitating

transcription (PubMed:30867594). Acts as a mediator of neurotransmission-independent role of nuclear dopamine in ventral tegmental area (VTA) neurons: catalyzes dopaminylation of 'Gln- 5' of histone H3 (H3Q5dop), thereby regulating relapse-related transcriptional plasticity in the reward system (PubMed:32273471). Regulates vein remodeling by mediating serotonylation and subsequent inactivation of ATP2A2/SERCA2 (By similarity). Also acts as a protein deamidase by mediating the side chain deamidation of specific glutamine residues of proteins to glutamate (PubMed:9623982, PubMed:20547769). Catalyzes specific deamidation of protein gliadin, a component of wheat gluten in the diet (PubMed:9623982). May also act as an isopeptidase cleaving the previously formed cross-links (PubMed:26250429, PubMed:27131890). Also able to participate in signaling pathways independently of its acyltransferase activity: acts as a signal transducer in alpha-1 adrenergic receptor-mediated stimulation of phospholipase C-delta (PLCD) activity and is required for coupling alpha-1 adrenergic agonists to the stimulation of phosphoinositide lipid metabolism (PubMed:8943303).

Cellular Location

Cytoplasm, cytosol. Nucleus. Chromosome. Secreted, extracellular space, extracellular matrix. Cell membrane {ECO:0000250|UniProtKB:Q9WVJ6}. Mitochondrion. Note=Mainly localizes to the cytosol (PubMed:9575137). Present at much lower level in the nucleus and chromatin (PubMed:9575137). Also secreted via a

non-classical secretion pathway to the
extracellular matrix (PubMed:27270573)

Goat Anti-TGM2 Antibody - Protocols

Provided below are standard protocols that you
may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)